

On the quantum cohomology of adjoint varieties

P.E. Chaput, N. Perrin

April 30, 2009

Abstract

We study the quantum cohomology of quasi-minuscule and quasi-cominuscule homogeneous spaces. The product of any two Schubert cells does not involve powers of the quantum parameter higher than 2. With the help of the quantum to classical principle we give presentations of the quantum cohomology algebras. These algebras are semi-simple for adjoint non coadjoint varieties and some properties of the induced strange duality are shown.

1 Introduction

In this paper, we study the quantum and classical cohomology rings of some rational homogeneous varieties. The papers [ChMaPe07], [ChMaPe08], [ChMaPe09a] dealt with the quantum cohomology of minuscule and cominuscule rational homogeneous spaces. Here we are interested in quasi-minuscule and quasi-cominuscule varieties (see Definition 2.1).

The choice of these varieties comes from the fact that Schubert calculus has a simpler and in many cases combinatorial expression: jeu de Taquin and its generalisations. This was remarked for (co)minuscule varieties by H. Thomas and A. Yong in [ThYo08] and generalised for a large class of Schubert classes (called Λ -minuscule, see Definition 2.9), in [ChPe09a]. In particular, for quasi-(co)minuscule varieties, all Schubert classes up to half the dimension are Λ -minuscule (see Corollary 2.12). This enables computing presentations of the classical cohomology ring of these varieties.

Moreover, as for the (co)minuscule case, we will explain here that the quantum Schubert calculus is also simpler for these varieties, and give several technics for computing Gromov-Witten invariants. The first one (see Theorem 2) is a simple interpretation of the quantum Chevalley formula as given in [FuWo04]. We then prove a result on the power of q , the quantum parameter, appearing in a product of two Schubert classes. In particular for X an adjoint of coadjoint variety (see Definition 2.1) we prove a vanishing result for the quantum Schubert calculus and relate it to the geometry of these varieties:

Theorem 1 *Let σ_u and σ_v be classical cohomology classes. Then if we write*

$$\sigma_u * \sigma_v = \sum_{d,w} c_{u,v}^w(d) q^d \sigma_w \quad ,$$

we have $c_{u,v}^w(d) = 0$ unless $d \leq 2$.

Key words: Quantum cohomology, adjoint varieties, strange involution.
 Mathematics Subject Classification: 14M15, 14N35

We also explain that the quantum to classical principle as initiated in [BuKrTa03] can also be applied to compute some Gromov-Witten invariants. With all these technics, we are able to give presentations of the quantum cohomology rings of all quasi-(co)minuscule homogeneous spaces. These results are presented in Section 5. We do not give here a presentation in the classical cases as they were obtained by A. Buch, A. Kresch and H. Tamvakis in [BuKrTa08] but focus on exceptional groups. Some of the computations were made using a computer program written in Java (and available at the webpage www.math.sciences.univ-nantes.fr/~chaput/quiver-demo.html), which may be of some interest for those who want to make explicit computations in these quantum cohomology rings.

Finally we focus on strange duality, a known phenomenon for (co)minuscule homogeneous spaces X : in that case there exists an involution ι of the quantum cohomology algebra $QH^*(X, \mathbb{C})_{loc}$ with the quantum parameter q inverted sending classes of degree d to classes of degree $-d$. This involution, in the (co)minuscule cases also sends a Schubert class to a multiple of a Schubert class. The existence of such an involution is a consequence of the semi-simplicity of the ring $QH^*(X, \mathbb{Z})_{loc}$. We conjectured in [ChMaPe09a] that the only homogeneous spaces X with $QH^*(X, \mathbb{Z})_{loc}$ semi-simple are the (co)minuscule homogeneous spaces.

In Section 6 we are able to prove, for many examples, that for X a rational homogeneous space with Picard number one the algebra $QH^*(X, \mathbb{Z})_{loc}$ is not semi-simple (see Theorem 4). However, we give counterexamples to our conjecture and prove that for X an adjoint non coadjoint variety, the localised algebra $QH^*(X, \mathbb{C})_{loc}$ is semi-simple (see Theorem 5). This implies the existence of an algebra involution ι as above. We prove that this involution, once specialised at $q = 1$, is the identity on classes of degree multiple of $c_1(X)$ (Proposition 6.7) but is more complicated in general than the involution in the minuscule and cominuscule cases (see Theorem 6). In particular, we are not able to compute it except for very special classes.

Contents

1	Introduction	1
2	Minuscule, Cominuscule, Adjoint and Coadjoint varieties	3
2.1	First definitions	3
2.2	Roots and Schubert varieties in the adjoint and coadjoint cases	4
2.3	Description of ϖ -minuscule elements in the adjoint and coadjoint cases	5
2.4	Quantum Hasse diagram	6
2.5	Affine symmetries in the adjoint and coadjoint cases	9
3	Maximal degree for quantum cohomology	9
3.1	Dimension argument	9
3.2	Rational curves on homogeneous spaces	10
3.3	Adjoint case	12
3.4	Coadjoint varieties as hyperplane sections of Scorza varieties	13
3.5	A description of Y'_3 for Scorza varieties	14
3.6	Y'_3 for a hyperplane section of a Scorza variety	17
4	Quantum to classical principle for degree one Gromov-Witten invariants	18
5	Presentation of the quantum cohomology rings	21
5.1	Adjoint and coadjoint varieties in type G_2	21
5.2	Adjoint and coadjoint varieties in type F_4	21

5.3	Adjoint and coadjoint varieties in type E_6	22
5.4	Adjoint and coadjoint varieties in type E_7	24
5.5	Adjoint and coadjoint varieties in type E_8	26
6	Semi-simplicity of the quantum cohomology and strange duality	27
6.1	Coadjoint varieties	28
6.2	Adjoint non coadjoint varieties	29
6.3	Some properties of the involution for adjoint non coadjoint varieties	32
7	The incidence variety	36

2 Minusculer, Cominusculer, Adjoint and Coadjoint varieties

Let G be a semi-simple algebraic group. Let T be a maximal torus in G and B a Borel subgroup containing T . We denote by W the associated Weyl group, with Δ the root system and with S the set of simple roots. We shall denote by Θ the highest root and with θ the highest short root. For simply-laced groups, we have $\Theta = \theta$. We will denote by α_{ad} any simple root such that $\langle \alpha_{ad}, \Theta^\vee \rangle = 1$ (such a root is unique except in type A).

2.1 First definitions

Definition 2.1 Let ϖ be a dominant weight, we shall call ϖ

- *minuscule* if for any positive root $\alpha \in R_+$ we have $\langle \alpha^\vee, \varpi \rangle \leq 1$;
- *cominuscule* if it is minuscule for the dual root system ;
- *quasi-minuscule* if for any positive root $\alpha \in R_+$ we have $\langle \alpha^\vee, \varpi \rangle \leq 2$ with equality iff $\alpha = \varpi$;
- *quasi-cominuscule* if it is quasi-minuscule for the dual root system.

Remark 2.2 A non minuscule quasi-minuscule weight ϖ is a root. More precisely we have $\varpi = \theta$. A quasi-cominuscule non cominuscule weight ϖ is also a root and we have $\varpi = \Theta$. In this case we shall say that ϖ is *adjoint*. We call a weight *coadjoint* if it is adjoint for the dual root system, in this case we have $\varpi = \theta$.

In simply laced cases, minuscule and cominuscule weights, quasi-minuscule and quasi-cominuscule weights and adjoint and coadjoint weights coincide.

Type of G	Quasi-minuscule		Quasi-cominuscule	
	Minuscule	Coadjoint	Cominuscule	Adjoint
A_n	$\varpi_1, \dots, \varpi_n$	$\varpi_1 + \varpi_n$	$\varpi_1, \dots, \varpi_n$	$\varpi_1 + \varpi_n$
B_n	ϖ_n	ϖ_1	ϖ_1	ϖ_2
C_n	ϖ_1	ϖ_2	ϖ_n	ϖ_1
D_n	$\varpi_1, \varpi_{n-1}, \varpi_n$	ϖ_2	$\varpi_1, \varpi_{n-1}, \varpi_n$	ϖ_2
E_6	ϖ_1, ϖ_6	ϖ_2	ϖ_1, ϖ_6	ϖ_2
E_7	ϖ_7	ϖ_1	ϖ_7	ϖ_1
E_8	None	ϖ_8	None	ϖ_8
F_4	None	ϖ_4	None	ϖ_1
G_2	None	ϖ_1	None	ϖ_2

Minuscule, Quasi-minuscule, Cominusculer, Quasi-cominusculer, Adjoint and Coadjoint weights.

Definition 2.3 Let P be the standard parabolic subgroup associated to a weight ϖ . We will say that the homogeneous variety G/P is *minuscule*, *cominuscule*, *quasi-minuscule*, *quasi-cominuscule*, *adjoint*

or *coadjoint* if the weight ϖ is minuscule, cominuscule, quasi-minuscule, quasi-cominuscule, adjoint or coadjoint.

Remark 2.4 (i) The adjoint homogeneous spaces correspond to the classical terminology of adjoint varieties.

(ii) The geometry of minuscule homogeneous varieties is better understood than the geometry of general homogeneous spaces. See for example [Ses78], [ThYo08] or [Per07]. Quasi-minuscule weights were in particular introduced to generalise the classical theory of standard monomials in minuscule varieties (see [Ses78]) to other homogeneous spaces (see [LaMuSe79]).

Remark 2.5 For minuscule and cominuscule homogeneous spaces, the quantum cohomology was described in [BuKrTa03] for classical groups and in [ChMaPe08] in general. So we will mainly focus in the sequel on adjoint and coadjoint homogeneous spaces except in Subsection 2.4 where we include the minuscule and cominuscule cases in our simplified version of quantum Chevalley formula.

The adjoint and coadjoint varieties are tabulated in the following array (for the simply-laced Lie algebras the notions of adjoint and coadjoint varieties coincide and therefore we leave the corresponding box empty):

	Adjoint				Coadjoint			
G	Θ	G/P_Θ	$\dim(G/P_\Theta)$	$c_1(G/P_\Theta)$	θ	G/P_θ	$\dim(G/P_\theta)$	$c_1(G/P_\theta)$
A_n	$\varpi_1 + \varpi_n$	$\mathbb{F}(1, n, n+1)$	$2n+1$	$(n+1, n+1)$				
B_n	ϖ_2	$\mathbb{G}_Q(2, 2n+1)$	$4n-5$	$2n-2$	ϖ_1	$\mathbb{Q}^{2n-1}$	$2n-1$	$2n-1$
C_n	$2\varpi_1$	$\mathbb{P}^{2n-1}$	$2n-1$	n	ϖ_2	$\mathbb{G}_\omega(2, 2n)$	$4n-5$	$2n-1$
D_n	ϖ_2	$\mathbb{G}_Q(2, 2n)$	$4n-7$	$2n-3$				
E_6	ϖ_2		21	11				
E_7	ϖ_1		33	17				
E_8	ϖ_8		57	29				
F_4	ϖ_1		15	8	ϖ_4		15	11
G_2	ϖ_2		5	3	ϖ_1	$\mathbb{Q}^5$	5	5

Invariants of Adjoint and Coadjoint varieties. (1)

In this array, we denote by $\mathbb{F}(1, n, n+1)$, $\mathbb{G}_Q(2, n)$, $\mathbb{G}_\omega(2, 2n)$ resp. $\mathbb{Q}^n$ the point-hyperplane incidence in $\mathbb{P}^n$, the Grassmannian of 2 dimensional isotropic subspaces in an n -dimensional subspace endowed with a nondegenerate quadratic form, the Grassmannian of 2 dimensional isotropic subspaces in an $2n$ -dimensional subspace endowed with a symplectic form resp. a n -dimensional quadric. We denote by $c_1(X)$ the index of the variety X . In the case of type C_n , the adjoint variety is isomorphic to $\mathbb{P}^{2n-1}$, but the projective embedding is given by twice the generator of $\text{Pic}(\mathbb{P}^{2n-1})$. Note that we always have $\dim(G/P_\Theta) = 2c_1(G/P_\Theta) - 1$ (this well-known fact can be seen as a consequence of Propositions 3.11 and 3.8).

2.2 Roots and Schubert varieties in the adjoint and coadjoint cases

In this subsection, we recall some results on Schubert varieties specific to the adjoint and coadjoint homogeneous spaces. We shall need the following:

Notation 2.6 (i) We define the affine weight ρ by $\langle \rho, \alpha^\vee \rangle = 1$ and the affine coweight $\rho^\vee$ by $\langle \alpha, \rho^\vee \rangle = 1$ for all simple root α of the affine Weyl group.

(ii) Recall that there is a natural order on the roots defined by $\alpha \leq \beta$ if $\beta - \alpha$ is in the monoid generated by positive roots.

Fact 2.7 Let $X = G/P$ be a coadjoint (respectively an adjoint) homogeneous space. Then, the Schubert varieties are in one to one correspondence with short (respectively long) roots. The corresponding map from W^P to this set of roots is given by $u \mapsto \alpha = u(\varpi)$ and we set $\sigma_\alpha = \sigma_u$.

Proof. This is a reformulation of the fact that the orbit of the point ϖ in X under the action of the Weyl group is the set of T -fixed points in X . $\square$

Let $u \in W^P$ and $\alpha = u(\varpi)$. We denote by $X(\alpha)$ the Schubert variety such that $[X(\alpha)] = \sigma_\alpha$ in $H^{2l(u)}(X, \mathbb{Z})$.

Proposition 2.8 Let $X = G/P$ be a coadjoint (respectively an adjoint) homogeneous space and let α be a short (respectively long) root. Denote by $X(\alpha)$ the associated Schubert variety. Then, the following propositions hold:

- (i) The integer $\dim(X)$ is odd, we define r by the equality $\dim(X) = 2r + 1$.
- (ii) There is an equivalence

$$X(\alpha) \subset X(\beta) \Leftrightarrow \begin{cases} \alpha \leq \beta & \text{for } \alpha \text{ and } \beta \text{ of the same sign,} \\ \text{Supp}(\alpha) \cup \text{Supp}(\beta) \text{ is connected} & \text{for } \alpha \text{ negative and } \beta \text{ positive.} \end{cases}$$

(iii) Denote by $X(\alpha)^\vee$ the Poincaré dual of $X(\alpha)$ and by i the Weyl involution, then we have the equality $X(\alpha)^\vee = X(-i(\alpha))$.

Proof. All these results can be easily deduced from results in [LaMuSe79, Section 3.1] in the coadjoint case. Indeed, (i) follows from Theorem 3.1 in *loc. cit.*, (ii) follows from the conjunction of Theorem 3.1 and Theorem 3.10 in *loc. cit.*. For (iii), recall that Poincaré duality is given on W^P by $u \mapsto w_0 u w_0 w_X$ where w_0 is the longest element in W and w_X the longest element in W^P . In terms of roots, if $\alpha = u(-\varpi)$, then the root corresponding to $X(\alpha)^\vee$ is

$$w_0 u w_0 w_X(-\varpi) = w_0 u w_0(\varpi) = w_0 u(-\varpi) = w_0(\alpha) = -i(\alpha).$$

To prove the corresponding statements in the adjoint case, little more is needed. Indeed, all the results only involve Weyl groups and are therefore true simply because the Weyl group of a root system and the Weyl group of its dual are isomorphic. $\square$

2.3 Description of ϖ -minuscule elements in the adjoint and coadjoint cases

Let us recall, according to Dale Peterson [Pro99a, p.273] (see also [ChPe09a]), the following definition:

Definition 2.9 Let $\Lambda = \sum_i \Lambda_i \varpi_i$ be a dominant weight (where $(\varpi_i)_i$ are the fundamental weights).

- An element $w \in W$ is Λ -minuscule if there exists a reduced decomposition $w = s_{i_1} \cdots s_{i_l}$ such that for any $k \in [1, l]$ we have $s_{i_k} s_{i_{k+1}} \cdots s_{i_l}(\Lambda) = s_{i_{k+1}} \cdots s_{i_l}(\Lambda) - \alpha_{i_k}$.
- w is Λ -cominuscule if w is $(\sum \Lambda_i \varpi_i^\vee)$ -minuscule.

Remark 2.10 For P a parabolic subgroup associated to a (co)minuscule weight ϖ , then any element in W^P is ϖ -(co)minuscule.

Let $X = G/P$ be a coadjoint or an adjoint variety associated to the fundamental weight ϖ . Recall that there exists a non negative integer r such that $\dim(X) = 2r + 1$.

Proposition 2.11 *If X is coadjoint (resp. adjoint), then any element w in W^P with $l(w) \leq r$ is ϖ -minuscule (resp. ϖ -cominuscule).*

Proof. The proof of Theorem 3.1 in [LaMuSe79] proves that, in the coadjoint case, such a u is ϖ -minuscule and that it is maximal for this property. The adjoint case is obtained from the previous one by taking the dual root system. $\square$

We refer to [ChPe09a] for a definition of jeu de taquin. As a consequence of the main result in *loc. cit.* we have:

Corollary 2.12 *Let u, v and w be elements in W^P , then the Littlewood-Richardson coefficient $c_{u,v}^w$ can be computed using jeu de taquin for $l(w) \leq r$.*

2.4 Quantum Hasse diagram

In this subsection, we shall give a simplified statement for the quantum Chevalley formula as formulated in [FuWo04]. We shall also see that our parametrisation of Schubert classes by roots as in the previous subsection extends to the quantum monomials, which are by definition of the form $q^d \sigma_u$ for some integer d and some $u \in W^P$: these monomials are parametrised by the roots of the affine root system $\hat{R}$ associated to R and this parametrisation is compatible with the quantum Chevalley formula.

For the moment we only assume that X has Picard number one. Let us first define an injection of the set of quantum monomials in the set of affine weights P^{aff} .

Definition 2.13

- We denote $\mathcal{M}$ the set of quantum monomials (with q inverted), namely the set of pairs (w, d) in $W^P \times \mathbb{Z}$.
- Let $w \in W^P$ and $d \in \mathbb{Z}$. To the quantum monomial $\sigma_w \cdot q^d$ we associate the affine weight $\eta(w, d) = u(\varpi) - d\delta$, where δ is the minimal positive imaginary root in $\hat{R}$ (see for example [Kac90]).
- For $\pi = \eta(w, d)$ we denote $\sigma_\pi = \sigma_w \cdot q^d$ the corresponding quantum monomial.
- For $\pi \in \eta(\mathcal{M})$ we denote $l(\pi)$ the integer $l(w) + dc_1$, if (w, d) is the element such that $\eta(\sigma_w, d) = \pi$.

Remark 2.14 In the coadjoint case, $\eta(\mathcal{M})$ is the set of all short real roots while in the adjoint case it is the set of all long real roots.

Proposition 2.15 *Let $\pi \in \eta(\mathcal{M})$; we have*

$$h * \sigma_\pi = \sum_{\gamma > 0, l(s_\gamma(\pi)) = l(\pi) + 1} \langle \pi, \gamma^\vee \rangle \sigma_{s_\gamma(\pi)}.$$

Proof. This is the particular case, when the Picard number of X is one, of [Ful97, Theorem 10.1]. In fact, let us assume that $\pi = \eta(w, 0)$ with $w \in W^P$; according to this result, the quantum product of h and σ_w is

$$\sum \langle \varpi, \alpha^\vee \rangle \sigma_{us_\alpha} + \sum q^{d(\alpha)} \langle \varpi, \alpha^\vee \rangle \sigma_{us_\alpha} \quad (2)$$

where α is a positive root, $d(\alpha) = \langle \varpi, \alpha^\vee \rangle$, and in the first resp. second summand we have $l(ws_\alpha) = l(w) + 1$ resp. $l(ws_\alpha) = l(w) + 1 - c_1 d(\alpha)$. We denote A resp. A_1, A_2 the set of these roots resp. the

roots in the first, second case. We consider an injection $\varphi : A \rightarrow \widehat{R}$ by mapping $\alpha \in A_1$ to $w(\alpha)$ and $\alpha \in A_2$ to $w(\alpha) + \delta$.

Let $\alpha \in A$ and let us denote $\gamma = \varphi(\alpha)$. We claim that for $\alpha \in A_1$ resp. $\alpha \in A_2$, $s_\gamma w(\varpi)$ is equal to $ws_\alpha(\varpi)$ resp. $ws_\alpha(\varpi) - d(\alpha)\delta$. In fact the first case follows from the equality $ws_\alpha = ws_\alpha w^{-1}w = s_{w(\alpha)}w$. The second case may be computed as follows:

$$\begin{aligned} s_\gamma w(\varpi) &= s_{w(\alpha)+\delta}(\varpi) = w(\varpi) - \langle w(\alpha)^\vee, w(\varpi) \rangle (w(\alpha) + \delta) \\ &= w(\varpi) - \langle \alpha^\vee, \varpi \rangle w(\alpha) - d(\alpha)\delta = ws_\alpha(\varpi) - d(\alpha)\delta, \end{aligned}$$

where the last equality follows from the first case. Thus in particular we get $s_\gamma w(\varpi) = \eta(ws_\alpha, d(\alpha))$.

Moreover it follows that γ is a positive root. In fact, in the second case it has positive coefficient on δ , and in the first case we have $s_\gamma w(\varpi) = w(\varpi) - d(\alpha)\gamma$ and on the other hand $s_\gamma w(\varpi) = ws_\alpha(\pi) < w(\alpha)$, so that $\gamma > 0$.

Finally φ is a bijection from A to the set of positive roots γ such that $l(s_\gamma w(\pi)) = l(w(\pi)) + 1$, and the proposition is proved. $\square$

Our next goal is to restrict the set of root γ in the former proposition in the quasi-(co)minuscule cases. We express the index c_1 of our varieties in terms of the highest roots. We start with a:

Lemma 2.16 *Let $X = G/P$ be a homogeneous space corresponding to a fundamental weight ϖ and let β be the simple root with $\langle \varpi, \beta^\vee \rangle = 1$. Let $\alpha^\vee$ be the biggest coroot with the following properties:*

- $\langle \varpi, \alpha^\vee \rangle = 1$.
- $|\alpha^\vee| = |\beta^\vee|$.

Then we have $c_1(X) = \langle \rho, \alpha^\vee \rangle + 1$.

Proof. Let us recall (see [BrKu05, Page 85]) that $c_1(X) = \langle \delta_P, \beta^\vee \rangle$ where $\delta_P = \rho + w^P \rho$ and w^P is the longest element in W_P the Weyl group of P . To prove the lemma it suffices to explain why $w^P(\beta^\vee) = \alpha^\vee$. The coroot $w^P(\beta^\vee)$ must have the same length as $\beta^\vee$.

First let us prove that the set B of coroots $\alpha^\vee$ of the same length as $\beta^\vee$ and with $\langle \varpi, \alpha^\vee \rangle = 1$ is equal to the orbit of $\beta^\vee$ under W_P . Clearly the orbit is contained in B . Now let $\beta_0^\vee \in B$. We proceed by induction on $\langle \beta_0^\vee, \rho \rangle$. If $\langle \beta_0^\vee, \rho \rangle = 1$, then $\beta_0^\vee = \beta^\vee$ and we are done. Otherwise, there exists $\gamma^\vee$ a simple coroot of P with $\langle \gamma, \beta_0^\vee \rangle > 0$. Indeed, otherwise, we would have $\langle \rho_P, \beta_0^\vee \rangle \leq 0$ where ρ_P is half the sum of the roots of the Levi part $L(P)$ of P . But because $\beta_0^\vee$ is positive this would imply $\langle \rho_P, \beta_0^\vee \rangle = 0$ thus $\beta_0^\vee = \beta^\vee$. Now $s_{\gamma^\vee}(\beta_0^\vee)$ is again in B with $\langle \rho, s_{\gamma^\vee}(\beta_0^\vee) \rangle < \langle \rho, \beta_0^\vee \rangle$ and the result on B follows by induction.

We end by proving that $w^P(\beta^\vee)$ is the highest coroot of B . Indeed, if u and v are in W_P with $u \leq v$, and if ϖ' is a fundamental weight of $L(P)$, then $u(\varpi') \geq v(\varpi')$ i.e. $u(\varpi') - v(\varpi')$ is a sum of positive roots of $L(P)$. We get $\langle u(\varpi') - v(\varpi'), \beta^\vee \rangle \leq 0$ thus $\langle u(\beta), \varpi'^\vee \rangle \leq \langle v(\alpha), \varpi'^\vee \rangle$ and $u(\beta^\vee) \leq v(\beta^\vee)$. $\square$

If R is a root system, remark that the coroot $\Theta^\vee$ is the highest short root of the dual root system while $\theta^\vee$ is the highest root of the dual root system. From now on we assume that X is quasi-(co)minuscule and has Picard rank one. We have:

Lemma 2.17 *The following formulas yield the first Chern class of minuscule, cominuscule, coadjoint or adjoint varieties:*

	<i>Minuscule</i>	<i>Coadjoint</i>	<i>Cominuscule</i>	<i>Adjoint</i>
c_1	$\langle \rho, \theta^\vee \rangle + 1$	$\langle \rho, \theta^\vee \rangle$	$\langle \rho, \Theta^\vee \rangle + 1$	$\langle \rho, \Theta^\vee \rangle$
c_1	$\langle \Theta, \rho^\vee \rangle + 1$	$\langle \Theta, \rho^\vee \rangle$		

Proof. If G is of type A , then X is a Grassmannian and the lemma is true. If not, let $\alpha_{ad}^\vee$ denote the adjoint simple coroot, that is the only coroot such that $\langle \alpha_{ad}, \Theta^\vee \rangle = 1$. To prove the first line it is enough to check that the root α of Lemma 2.16 is $\theta^\vee, \theta^\vee - \alpha_{ad}^\vee, \Theta^\vee, \Theta^\vee - \alpha_{ad}^\vee$, according to the four cases.

In the minuscule case, the simple root β with $\langle \varpi, \beta^\vee \rangle = 1$ is short and therefore $\beta^\vee$ is long; moreover by definition of a minuscule weight we have $\langle \varpi, \gamma^\vee \rangle \leq 1$ for any coroot $\gamma^\vee$. It thus follows that $\alpha^\vee$ is the highest coroot.

In the cominuscule case, $\beta^\vee$ must be short. Moreover for any short coroot $\gamma^\vee$ we have $\langle \varpi, \gamma^\vee \rangle \leq 1$. Thus in this case $\alpha^\vee = \Theta^\vee$.

The coadjoint and adjoint cases are similar, except that the highest coroot $\gamma^\vee$ of the same length as $\beta^\vee$ has coefficient 2 on $\beta^\vee$, so that we must have $\alpha^\vee = \gamma^\vee - \alpha_{ad}^\vee$.

To prove the second line of the table, we only need to apply the quasi-cominuscule case to the dual root system. We may also observe that the sum of the coefficients of the highest root is the same in a root system and its dual (for the minuscule and coadjoint cases). $\square$

Lemma 2.18 *If $[a/b]$ denotes the integral part of a/b , we have the following equalities:*

	<i>Minuscule</i>	<i>Coadjoint</i>
$\langle \varpi - \pi, \rho^\vee \rangle$	$l(\pi)$	$l(\pi) + [l(\pi)/c_1]$
	<i>Cominuscule</i>	<i>Adjoint</i>
$\langle \rho, (\varpi - \pi)^\vee \rangle$	$l(\pi)$	$l(\pi) + [l(\pi)/c_1]$

Proof. Note that we have $\langle \delta, \rho^\vee \rangle = \langle \Theta, \rho^\vee \rangle + 1$ since $\delta = \Theta + \alpha_0$. Thus using Lemma 2.17 and the definition of $l(\pi)$ given in Definition 2.13, it is enough to prove the lemma when $\pi = \eta(w, 0)$.

In this case, assuming we are in the minuscule case, if we write $w = s_{i_l} \cdots s_{i_1}$, we have $w(\varpi) = \varpi - \sum \alpha_{i_k}$, and so $\langle \varpi - w(\varpi), \rho^\vee \rangle = l = l(w)$. Thus the lemma is proved in this case. The cominuscule case follows by duality.

In the coadjoint case, if $l(w) < c_1$, then any reduced expression of w will have the same property as in the minuscule case and the same argument gives $\langle \varpi - w(\varpi), \rho^\vee \rangle = l(w)$. If $l(w) \geq c_1$, then we can find a reduced expression $w = s_{i_l} \cdots s_{i_1}$ such that $w(\varpi) = \varpi - \sum d_k \alpha_{i_k}$ where $d_k = 1$ for all k but $k = c_1$, and $d_{c_1} = 2$. Thus $\langle \varpi - w(\varpi), \rho^\vee \rangle = l(w) + 1$ and the lemma is again true. The adjoint case follows by duality. $\square$

Definition 2.19 Let $\pi \in \eta(\mathcal{M})$ and γ be a root. We say that γ interacts with π if $\langle \pi, \gamma^\vee \rangle > 0$ and either:

- We are in the minuscule or cominuscule cases and γ is a simple root.
- We are in the coadjoint or adjoint cases, c_1 does not divide $l(\pi) + 1$, and γ is a simple root.
- We are in the coadjoint or adjoint cases and c_1 divides $l(\pi) + 1$; this implies that there exist a simple root α and an integer d such that $\pi = \alpha + d\delta$. In this case γ interacts with π if either $\gamma = \alpha$ or $\gamma = \alpha + \beta$, where β is another simple root.

We are now in a position to give a more explicit formula for the quantum Chevalley formula of Proposition 2.15:

Theorem 2 *We have a formula*

$$h * \sigma_\pi = \sum_{\gamma} \langle \pi, \gamma^\vee \rangle \sigma_{s_\gamma(\pi)} ,$$

where the sum runs over all positive roots γ which interact with π .

Proof. Assume first we are in the minuscule case. Let $\gamma > 0$ be such that $l(s_\gamma(\pi)) = l(\pi) + 1$. Then by Lemma 2.18 we have $\langle \varpi - s_\gamma(\pi), \rho^\vee \rangle = \langle \varpi - \pi + \gamma, \rho^\vee \rangle = l(\pi) + 1 = \langle \varpi - \pi, \rho^\vee \rangle + 1$, thus $\langle \gamma, \rho^\vee \rangle = 1$ and γ is a simple root. In the cominuscule case the same argument gives that $\gamma^\vee$ is a simple root, so γ is simple. The coadjoint and adjoint cases are similar, except that one has to take care whether $l(s_\gamma(\pi))$ is divisible by c_1 or not. $\square$

2.5 Affine symmetries in the adjoint and coadjoint cases

The description of the quantum monomials in terms of roots gives a nice interpretation of the affine symmetries described in [ChMaPe09b]. Let us recall the description of these symmetries, we refer to [ChMaPe09b] for more details. Let c be a cominuscule vertex of the Dynkin diagram and let α_c be the corresponding cominuscule simple root. Denote by τ_c the orientation preserving automorphism of the extended Dynkin diagram sending the added vertex to the vertex c (there is a unique such automorphism). Denote by $\varpi_c^\vee$ the cominuscule coweight associated to c and by v_c the shortest element such that $v_c \varpi_c^\vee = w_0 \varpi_c^\vee$ with w_0 the longest element of the Weyl group. We proved in [ChMaPe09b] the following formula

$$\sigma_{v_c} * \sigma_u = q^{\eta_P(\varpi_c^\vee - u^{-1}(\varpi_c^\vee))} \sigma_{v_c u}$$

where $\eta_P : Q^\vee \rightarrow Q^\vee / Q_P^\vee$ is the projection from the coroot lattice to its quotient by the coroot lattice of P and where we identify this quotient with $\mathbb{Z}$.

Proposition 2.20 (i) *We have the equality*

$$\sigma_{v_c} = \begin{cases} \sigma_{-\alpha_c - \Theta + \theta} & \text{in the coadjoint case} \\ \sigma_{-\alpha_c} & \text{in the adjoint case.} \end{cases}$$

In particular $\deg(\sigma_{v_c}) = c_1$.

(ii) *For any real long root α in the adjoint case (resp. short root α in the coadjoint case), we have the formula*

$$\sigma_{v_c} * \sigma_\alpha = \sigma_{\tau_c(\alpha) - \delta} = q \sigma_{\tau_c(\alpha)}.$$

Proof. Point (i) is an easy computation. For (ii), let $u \in W^P$ such that $u(\varpi) = \alpha$. Using the fact that $\tau_c = v_c t_{-\varpi_c^\vee}$ we get $\tau_c(\alpha) = v_c(\alpha) + \langle \alpha, \varpi_c^\vee \rangle \delta$. But we check the equality $\eta_P(\varpi_c^\vee - u^{-1}(\varpi_c^\vee)) = 1 - \langle \alpha, \varpi_c^\vee \rangle$ and the result follows. $\square$

3 Maximal degree for quantum cohomology

3.1 Dimension argument

In this section we want to bound the possible degrees of the quantum monomials in the quantum parameter q that may appear in a quantum product $\sigma_u * \sigma_v$. The first easy restriction is given by the following:

Lemma 3.1 *Let X be a adjoint or coadjoint homogeneous space. Let u and v in W^P , then if we write*

$$\sigma_u * \sigma_v = \sum_{d,w} c_{u,v}^w(d) q^d \sigma_w$$

we have $c_{u,v}^w(d) = 0$ unless $d \leq 3$.

Proof. This is a simple dimension count. Recall that $c_{u,v}^w(d) = 0$ unless $l(u) + l(v) + \dim X - l(w) = \dim X + c_1(X)d$ thus unless $c_1(X)d \leq 2 \dim X$. But in the adjoint case, we have $\dim X = 2c_1(X) - 1$ giving $c_{u,v}^w(d) = 0$ unless $d \leq 3$. In the coadjoint non adjoint case, there are three cases. For the Lagrangian Grassmannian, we have $\dim \mathbb{G}_\omega(2, 2n) = 4n - 5 = 2c_1(\mathbb{G}_\omega(2, 2n)) - 3$. Thus we deduce that $c_{u,v}^w(d) = 0$ if $d > 3$. For F_4/P_4 we have $\dim(F_4/P_4) = 15$ and $c_1(F_4/P_4) = 11$, thus $c_{u,v}^w(d) = 0$ unless $d \leq 2$. For G_2/P_1 , this variety being a quadric, the result is well known (see for example [ChMaPe07]). $\square$

In the next subsections we show that $c_{u,v}^w(d) = 0$ if $d > 2$. In the coadjoint non adjoint case, we are in fact more interested in understanding the geometric reason of this vanishing, since the proof of the above Lemma almost proves this point (the only case left is that of $\mathbb{G}_\omega(2, 2n)$). Rather what we will do is to describe in a uniform way the locus Y_3 covered by degree 3 curves passing through 2 general points (see Definition 3.2 below) for adjoint and coadjoint varieties and explain that the vanishing of the quantum coefficients is related to a porism. We believe that it is worth it describing the rich geometry of the subvarieties Y_3 .

3.2 Rational curves on homogeneous spaces

For the following definition, we shall assume that X is a homogeneous space with Picard group $\text{Pic}(X) = \mathbb{Z}$ and we pick the only ample line bundle $\mathcal{L}$ which is a generator of the Picard group (we may also include the adjoint variety for SL_n by taking as ample line bundle $\mathcal{L}$ the line bundle given by its minimal embedding). All degrees are considered with respect to $\mathcal{L}$.

Definition 3.2 Let d be an integer and x and y be points in X .

(i) Let $X_d(x)$ be the locus of points in X connected to x by a rational curve of degree d . In symbols: $X_d(x) = \{z \in X \mid \text{there exists } C \text{ a rational degree } d \text{ curve through } x \text{ with } z \in C\}$.

(ii) Let $Y_d(x, y)$ be the locus of points in X connected to x and y by the same rational curve of degree d . In symbols: $Y_d(x, y) = \{z \in X \mid \text{there exists } C \text{ a rational degree } d \text{ curve with } x, y, z \in C\}$.

We observed in [ChMaPe08] that, for X a minuscule or cominuscule rational homogeneous space, the geometry of rational curves and especially the geometry of the varieties $X_d(x)$ and $Y_d(x, y)$ is intimately related to the quantum cohomology of X . We shall see that this is still the case for adjoint and coadjoint varieties.

Let us denote by $V = V(\varpi)$ the irreducible representation of highest weight ϖ . For π a weight of the representation V , we denote by x_π a vector of weight π . All points in X are equivalent and we therefore will often take x_ϖ , a highest weight vector of V , as general point in X (recall that X is the orbit of $[x_\varpi]$, the class of x_ϖ in $\mathbb{P}V$).

We introduce the notation $M_{d,3}(X)$ to denote the moduli space of degree d morphism from $\mathbb{P}^1$ to X with three marked points on $\mathbb{P}^1$. Recall from [Tho98] or [Per02] that $M_{d,3}(X)$ is irreducible smooth and has the expected dimension $\dim(X) + dc_1(X)$. In particular, we may speak of general degree d rational curves on X . We shall denote by $ev_i : M_{d,3}(X) \rightarrow X$ the evaluation map defined by the i -th marked point.

Proposition 3.3 *Let x, y be two generic points on a generic curve of degree d , then $X_d(x)$ and $Y_d(x, y)$ are irreducible.*

Proof. We know that $M_{d,3}(X)$ is irreducible. We have a G -equivariant evaluation $ev_1 : M_{d,3}(X) \rightarrow X$. As X is covered by rational curves of any degree, its image is X itself. Assume that $x = x_\varpi$. Let U be a unipotent subgroup of G such that $U \rightarrow X$ is an isomorphism from U to an open neighbourhood of x , which we denote by O . We have an isomorphism

$$\begin{aligned} ev_1^{-1}(x_\varpi) \times U &\rightarrow ev_1^{-1}(O) \\ (f, u) &\mapsto u \circ f \end{aligned}$$

Since $ev_1^{-1}(O)$, an open subset of $M_{d,3}(X)$, is irreducible, $ev_1^{-1}(x)$ must be also irreducible. Therefore its image under ev_2 which is $X_d(x_\varpi)$, is irreducible.

For $Y_d(x, y)$ we argue essentially in the same way. We may assume that $x = x_\varpi$. We note that $X_d(x_\varpi)$ is a Schubert variety since it is irreducible from the above and B -stable since x_ϖ is fixed by B . Because y and the degree d curve through x_ϖ are supposed to be general, we may assume that y is in the open B -orbit of $X_d(x_\varpi)$. Thus there is again a unipotent subgroup U of B such that $U \rightarrow X$ induces an isomorphism between U and an open neighbourhood of y in $X_d(x_\varpi)$. Since U is included in B it preserves $ev_1^{-1}(x_\varpi)$, and thus induces an isomorphism $ev_1^{-1}(x_\varpi) \simeq (ev_1 \times ev_2)^{-1}(x_\varpi, y) \times U$. Therefore $(ev_1 \times ev_2)^{-1}(x_\varpi, y)$ is irreducible and we conclude as before. $\square$

Note that the proof shows that $Y_d(x, y)$ does not depend, up to isomorphism, on the choice of the two points x, y on a generic curve of degree d . Thus we can define the following:

Definition 3.4 Let $X_d(X)$ (resp. $Y_d(X)$) be the isomorphism class of the varieties $X_d(x)$ (resp. $Y_d(x, y)$) for one (resp. two) generic point(s) x (resp. x and y) on a generic degree d curve. When no confusion on the variety X can be made we simplify $X_d(X)$ and $Y_d(X)$ in X_d and Y_d .

Corollary 3.5 *The variety X_d in X is a Schubert variety stable under P .*

Definition 3.6 (i) We shall denote by $x_\varpi(d)$ the T -stable point in the open B -orbit of the Schubert variety $X_d(x_\varpi)$.

(ii) For d, n integers we denote by $\delta_X(d, n)$ the dimension of the space of degree d curves passing through n generic points of a generic curve of degree d in X .

Example 3.7 The value of $\delta_{\mathbb{P}^k}(1, n)$ is $2(k-1)$ if $n=0$, it is $k-1$ if $n=1$, and it is 0 otherwise.

We now prove a formula involving the dimension of $X_d(x)$ and of $Y_d(x, y)$.

Proposition 3.8 *Let x, y be two generic points on a generic degree d curve, then we have*

$$\dim(X_d(x)) + \dim(Y_d(x, y)) + \delta(d, 3) = c_1 d.$$

Proof. The moduli space $M_{d,3}(G/P)$ has dimension $\dim(X) + dc_1$. Therefore, as X is covered by rational curves of any degree, the inverse image $ev_1^{-1}(x_\varpi)$ has dimension dc_1 . Considering the morphism $ev_2 : ev_1^{-1}(x_\varpi) \rightarrow X$, with image $X_d(x_\varpi)$, we deduce the following equality of dimensions: $\dim((ev_1 \times ev_2)^{-1}(x_\varpi, x_\varpi(d))) = c_1 d - \dim(X_d)$.

Now the image of $ev_3 : (ev_1 \times ev_2)^{-1}(x_\varpi, x_\varpi(d)) \rightarrow X$ is by definition $Y_d(x, y)$, and the fibers of this morphism are $\delta(d, 3)$ -dimensional. $\square$

We recall here how to describe chains of T -invariant rational curves (see for example [FuWo04]):

Proposition 3.9 *The chains of k irreducible T -invariant rational curves starting from x_ϖ of degrees $(d_i)_{i \in [1,k]}$ are in bijection with sequences $(\alpha_i)_{i \in [1,k]}$ of positive roots with $\langle s_{\alpha_{m-1}} \cdots s_{\alpha_1}(\varpi), \alpha_m^\vee \rangle = d_m$ for all m . The end-point of the m -th component of such a chain is the projectivisation of the weight line of weight $s_{\alpha_m} \cdots s_{\alpha_1}(\varpi) = \varpi - d_1 \alpha_1 - \cdots - d_m \alpha_m$.*

We shall need the following:

Lemma 3.10 *Let $x, y \in X$ be two T -invariant points and d an integer. Assume that there is exactly one T -invariant chain of degree d through x and y . Then there is only one chain of degree d through x and y .*

Proof. Assume that there is a chain which is not T -invariant. Then the T -orbit of this chain provides a positive-dimensional family of chains of degree d through x and y . In the moduli space of curves, the corresponding positive dimensional variety must have at least 2 fixed points, a contraction. $\square$

3.3 Adjoint case

We first investigate the geometry of conics in adjoint homogeneous spaces so we fix an adjoint homogeneous space X .

Proposition 3.11 *We have $X_2 = X$ and Y_2 is a conic.*

Proof. Recall that Θ denotes the highest root. Since $\langle \Theta, \Theta \rangle = 2$, the curve through x_Θ with tangent direction $-\Theta$ has degree 2; its other T -fixed point has weight $s_\Theta(\Theta) = -\Theta$, thus it is $x_{-\Theta}$, that is, the T -fixed point representing the dense orbit. Therefore $X_2 = X$.

To understand Y_2 , we determine the degree 2 curves through x_Θ and $x_{-\Theta}$. By Proposition 3.9, such a curve is either given by a positive root α such that $\langle \Theta, \alpha \rangle = 2$ and $\Theta - 2\alpha = -\Theta$ (in which case $\alpha = \Theta$: this corresponds to the previous conic), or by a sequence (β, α) of two positive roots such that $\langle \Theta, \alpha \rangle = \langle s_\alpha(\Theta), \beta \rangle = 1$ and $\Theta - \alpha - \beta = -\Theta$. Since Θ is the highest root, this last equation implies $\alpha = \beta = \Theta$, contradicting the degrees.

Thus there is only one T -invariant degree 2 curve through x_Θ and $x_{-\Theta}$. By Lemma 3.10 there is only one degree 2 curve through x_Θ and $x_{-\Theta}$ and this implies that Y_2 is a conic. $\square$

Definition 3.12 We denote $Y'_3 \subset X$ the union of all the lines in X which meet Y_2 . It is also the union of all the *reducible* curves of degree 3 through two general points x and y .

Proposition 3.13 *Let x, y, z be three points in X . If there is a curve of degree 3 through x, y, z , then there are infinitely many. Thus $Y_3 = Y'_3$.*

Proof. Since the dimension of the space of curves through 3 points is an upper-continuous function on the 3 points, we may assume that x and y are generic, thus that $x = x_\Theta$ and $y = x_{-\Theta}$.

The proposition amounts to the fact that $\delta(3, 3) = 1$, or, using proposition 3.8, that $\dim Y_3 = c_1$. Let C be the conic through x and y and let Y'_3 be the variety covered by the lines meeting C . Since the dimension of the variety of lines through any point in X is $c_1 - 2$, the dimension of Y'_3 is at most c_1 . It is in fact c_1 since $T_x X$ and $T_y X$ only meet along the projectivisation of the line $[\mathfrak{g}_\Theta, \mathfrak{g}_{-\Theta}]$ in $\mathfrak{g}$.

To prove that $Y_3(x_\Theta, x_{-\Theta}) = Y'_3$, we use a suitable decomposition of $\mathfrak{g}$. Let $L \subset G$ be the pointwise stabiliser of C and $\mathfrak{l}$ its Lie algebra. If α is a root of $\mathfrak{l}$ then $\langle \Theta, \alpha^\vee \rangle = 0$. Let $\mathfrak{sl}_2(\Theta)$ be generated by $\mathfrak{g}_\Theta, \mathfrak{g}_{-\Theta}$ and $[\mathfrak{g}_\Theta, \mathfrak{g}_{-\Theta}]$. We consider $\mathfrak{g}$ as a module over $\mathfrak{sl}_2 \times \mathfrak{l}$. This module certainly contains $\mathfrak{sl}_2 \times \mathfrak{l}$. The tangent space $T_x X$ of X at x (resp. y) can be identified with the T -stable subspace

of $\mathfrak{g}$ with weights the negative roots α such that $\langle \Theta, \alpha^\vee \rangle < 0$ (resp. the positive roots α such that $\langle \Theta, \alpha^\vee \rangle > 0$). We define the subspace $T_x^1 X$ of $T_x X$ (resp. $T_x^1 X$ of $T_x X$) as the T -stable subspace with weights the negative roots α such that $\langle \Theta, \alpha^\vee \rangle = -1$ (resp. the positive roots α such that $\langle \Theta, \alpha^\vee \rangle = 1$). Since $L \subset P$, the space $V_L := T_x^1 X$ is a representation of L , included in $\mathfrak{g}$. Similarly $T_y^1 X$ is also an $\mathfrak{l}$ -representation, and moreover $T_x^1 X$ and $T_y^1 X$ are disjoint. Counting dimensions, we see that we have

$$\mathfrak{g} = \mathfrak{sl}_2 \times \mathfrak{l} \oplus \mathbb{C}^2 \otimes V_L$$

Concretely, here is the decomposition in each case :

$\mathfrak{g}$	Decomposition of $\mathfrak{g}$
$\mathfrak{sl}_n$	$(\mathfrak{gl}_2 \times \mathfrak{gl}_{n-2})_0 \oplus \mathbb{C}^2 \otimes (\mathbb{C}^{n-2} \oplus (\mathbb{C}^{n-2})^*)$
$\mathfrak{so}_{2n+1}$	$\mathfrak{sl}_2 \times (\mathfrak{sl}_2 \times \mathfrak{so}_{2n-3}) \oplus \mathbb{C}^2 \otimes (\mathbb{C}^2 \otimes \mathbb{C}^{2n-3})$
$\mathfrak{sp}_{2n}$	$\mathfrak{sl}_2 \times \mathfrak{sp}_{2n-2} \oplus \mathbb{C}^2 \otimes \mathbb{C}^{2n-2}$
$\mathfrak{so}_{2n}$	$\mathfrak{sl}_2 \times (\mathfrak{sl}_2 \times \mathfrak{so}_{2n-4}) \oplus \mathbb{C}^2 \otimes (\mathbb{C}^2 \otimes \mathbb{C}^{2n-4})$
$\mathfrak{e}_6$	$\mathfrak{sl}_2 \times \mathfrak{sl}_6 \oplus \mathbb{C}^2 \otimes \wedge^3 \mathbb{C}^6$
$\mathfrak{e}_7$	$\mathfrak{sl}_2 \times \mathfrak{so}_{12} \oplus \mathbb{C}^2 \otimes V^{32}$
$\mathfrak{e}_8$	$\mathfrak{sl}_2 \times \mathfrak{e}_7 \oplus \mathbb{C}^2 \otimes V^{56}$
$\mathfrak{f}_4$	$\mathfrak{sl}_2 \times \mathfrak{sp}_6 \oplus \mathbb{C}^2 \otimes \wedge_\omega^3 \mathbb{C}^6$
$\mathfrak{g}_2$	$\mathfrak{sl}_2 \times \mathfrak{sl}_2 \oplus \mathbb{C}^2 \otimes S^3 \mathbb{C}^2$

Using this description we can show that $Y_3 = Y'_3$. Let $\nu : \mathbb{C}^2 \rightarrow \mathfrak{sl}_2$ be the natural map of degree two with image the affine cone over $\mathfrak{sl}_2^{\text{ad}}$, the adjoint variety for $\mathfrak{sl}_2$. Let $X_L \subset V_L$ be the affine cone over the closed L -orbit in $\mathbb{P}V_L$. Then Y'_3 is described as the set of classes of elements $(s, 0, v \otimes \alpha) \in \mathfrak{sl}_2 \times \mathfrak{l} \oplus \mathbb{C}^2 \otimes V_L$, with $\alpha \in X_L$, and s and $\nu(v)$ proportional. The points x, y are respectively $[\nu(1, 0) : 0 : 0]$, $[\nu(0, 1) : 0 : 0]$.

If $z = [\nu(v), 0, v \otimes \alpha] \in Y'_3 \subset X$, $q : \mathbb{C}^2 \rightarrow \mathbb{C}$ is the quadratic form vanishing on $(1, 0)$ and $(0, 1)$ and such that $q(v) = 1$, l is a linear form such that $l(v) = 1$, then the map

$$\begin{aligned} \mathbb{P}^1 = \mathbb{P}\mathbb{C}^2 &\rightarrow Y'_3 \\ [w] &\mapsto [l(w). \nu(w) : 0 : q(w). w \otimes \alpha] \end{aligned}$$

is a rational curve of degree 3 through x, y and z . Thus there are infinitely many rational curves of degree 3 through x, y, z , so $Y'_3 = Y_3$ and the proposition is proved. $\square$

3.4 Coadjoint varieties as hyperplane sections of Scorza varieties

Scorza varieties were first defined and classified by F.L. Zak as varieties satisfying an extremal property with respect to their higher secants (see [Zak93]). For our purpose it will be enough to use the following classification theorem as a definition:

Definition 3.14 A Scorza variety is defined by two integers δ and m , with $\delta \in \{2, 4, 8\}$ and $m \geq 3$. If $\delta = 8$ then we must have $m = 3$. The corresponding homogeneous varieties are as follows (a generic hyperplane section of a Scorza variety turns out to be itself homogeneous; this homogeneous space is described in the third column):

δ	$X(\delta, m)$	$X(\delta) \cap H$
2	$\mathbb{P}^{m-1} \times \mathbb{P}^{m-1}$	$\mathbb{F}(1, m-1, m)$
4	$\mathbb{G}(2, 2m)$	$\mathbb{G}_\omega(2, 2m)$
8	E_6/P_1	F_4/P_4

The reason why we consider Scorza varieties is that we recover the coadjoint non adjoint varieties as hyperplane sections of some Scorza variety (except for G_2/P_1). Actually F.L. Zak's definition of Scorza varieties allows $\delta = 1$, but we don't need the varieties $X(1, m)$ and therefore will not speak about them.

Those varieties are naturally embedded via the smallest ample divisor $\mathcal{O}(1)$. The numbers δ, m have some geometrical and algebraic meanings: δ is the secant defect of $X(\delta, m)$, meaning that $\dim(\text{Sec}(X)) = 2\dim(X) + 1 - \delta$ where $\text{Sec}(X)$ denotes the union of the secant lines to X . More precisely through two generic points of X there passes a unique δ -dimensional quadric. Moreover m is the smallest integer such that the m -th higher secant of X is the whole projective space [Zak93].

On the other hand, the vector space in the projectivisation of which $X(\delta, m)$ is embedded has a natural structure of a Jordan algebra, the integer m is the generic rank of this algebra and δ the dimension of the composition algebra which coordinatises this algebra [Jac68].

Lemma 3.15 *We have $c_1(X) = \frac{\delta}{2}m$ and $\dim(X) = \delta(m - 1)$.*

Proof. The following numbers are well-known

X	$\mathbb{P}^{m-1} \times \mathbb{P}^{m-1}$	$\mathbb{G}(2, 2m)$	E_6/P_1
δ	2	4	8
$\dim(X)$	$2(m - 1)$	$2(2m - 2)$	16
$c_1(X)$	m	$2m$	12

and yield the lemma. □

3.5 A description of Y'_3 for Scorza varieties

In this subsection, we fix X a Scorza variety with parameters (δ, m) . We first describe Y_2 .

Proposition 3.16 *Through 2 generic points in X there passes a conic, and Y_2 is a smooth quadric of dimension δ .*

Proof. Let $x, y \in X$ be generic. By [Zak93], through x and y there passes a unique quadric of dimension δ , that we denote $Q_{x,y}$. This proves the first claim of the lemma. Moreover we have $Q_{x,y} \subset Y_2(x, y)$. Since $Y_2(x, y)$ is irreducible by Proposition 3.3, and has dimension at most $2c_1(X) - \dim(X) = \delta$ by Lemma 3.15 and Proposition 3.8, the proposition is proved. □

Definition 3.17 (i) Let $d > 0, e \geq 0$ be integers, Spin_{2d} the corresponding spinor group, A resp. $\mathbb{S}$ the vector resp. a spinor representation, with highest weight vector a_0 resp. s_0 . We denote $S \subset \mathbb{P}\mathbb{S}$ the spinor variety. Let W denote the natural representation of SL_e with highest weight vector w_0 .

(ii) We denote $S(d, e)$ the horospherical variety which is the closure of the $(\text{Spin}_{2d} \times SL_e)$ -orbit of $[a_0 + s_0 \otimes w_0]$ in $\mathbb{P}(A \oplus (S \otimes W))$ (for more on horospherical varieties, see for example [Pas08]). The closed orbit of $[a_0]$, which is a smooth quadric of dimension $2d - 2$, is a closed subvariety of $S(d, e)$ denoted $Q(d)$.

Lemma 3.18 *The variety $S(d, e)$ has dimension $\frac{(d+2)(d-1)}{2} + e$. Any point in $S(d, e) - Q(d)$ is contained in a linear space of dimension d included in $S(d, e)$, and intersecting $Q(d)$ along a maximal isotropic subspace (of dimension $d - 1$).*

Proof. Let $x = [a + s \otimes w]$ be a generic point in $S(d, e)$, with $a \in A, s \in \mathbb{S}, w \in W$ and $s \otimes w \neq 0$. Since x is in the $Spin_{2d} \times SL_e$ -orbit of $[a_0 + s_0 \otimes w_0]$, the class $[s]$ of s in $\mathbb{P}\mathbb{S}$ is a point of the spinor variety, thus it corresponds to a maximal isotropic linear subspace $L_d(s) \subset A$, and a is isotropic. Moreover $a \in L_d(s)$. Since the stabiliser in $Spin_{2d}$ of an element $[s] \in S$ acts transitively on the projectivisation of $L_d(s)$, $S(d, e)$ is the variety of all classes $[a' + s' \otimes w']$ satisfying $[s'] \in S$ and $a' \in L_d(s')$. It has dimension $d(d-1)/2 + (d-1) + (e-1) + 1$. Since $x \in S(d, e)$, $a \in L_d(s)$. Thus the linear space $\mathbb{P}(L_d(s) \oplus \mathbb{C} \cdot s \otimes w)$ satisfies the conditions of the lemma. $\square$

Recall that we denote by ϖ the dominant weight corresponding to X . For X a Scorza variety, remark that ϖ is minuscule. We define two subgroups of G as follows. Let P be the parabolic subgroup of G such that $G/P = X$ and denote by L its Levi subgroup. The Dynkin diagram of L can be recovered from the Dynkin diagram of G by removing the vertex v corresponding to P and the vertex $i(v)$ where i is the Weyl involution.

Definition 3.19 (i) We denote by L_1 the subgroup of L whose Dynkin diagram is the union of the connected components of the Dynkin diagram of L non orthogonal to Θ the highest root of G — or equivalently connected to the added vertex in the extended Dynkin diagram of G .

(ii) We denote by L_2 the subgroup of L whose Dynkin diagram is the union of the connected components of the Dynkin diagram of L orthogonal to Θ the highest root of G — or equivalently non connected to the added vertex in the extended Dynkin diagram of G .

(iii) We denote by $\tilde{L}_1$ the subgroup of G generated by L_1 and $SL_2(\Theta)$ (the subgroup of G whose Lie algebra is $\mathfrak{g}_{-\Theta} + [\mathfrak{g}_{-\Theta}, \mathfrak{g}_{\Theta}] + \mathfrak{g}_{\Theta}$). Its Dynkin diagram is the Dynkin diagram of L_1 together with the added vertex in the extended Dynkin diagram of G .

Pictures of the Dynkin diagrams of $G, L, \tilde{L}_1$ and L_2 for Scorza varieties are given in the proof of the following:




Proposition 3.20 *The variety $Y'_3(E_6/P_1)$ resp. $Y'_3(\mathbb{G}(2, 2m))$ is projectively equivalent to $S(5, 1)$ resp. $S(3, 2m-4)$. The variety $Y'_3(\mathbb{P}^{m-1} \times \mathbb{P}^{m-1})$ is a union of two irreducible components equivalent to $S(2, m-2) \simeq \mathbb{P}^1 \times \mathbb{P}^{m-1}$ and meeting along a smooth quadric surface.*

In all cases $Y'_3(X)$ is the linear section $X \cap \mathbb{P}(T_P \text{Sec}(X))$, where P is a generic point in the span of Y_2 .

Proof. Let α be a maximal root with vanishing pairing with $i(\varpi)$. In the case of E_6/P_1 resp. $\mathbb{G}(2, 2m)$, resp. $\mathbb{P}^{m-1} \times \mathbb{P}^{m-1}$, we have

$$\alpha = \begin{pmatrix} 1 & 2 & 2 & 1 & 0 \\ & & & 1 & \end{pmatrix} \text{ resp. } \alpha = (1, 1, \dots, 1, 0, 0), \text{ resp. } \alpha = \begin{cases} ((1, \dots, 1, 0), (0, \dots, 0)) \text{ or} \\ ((0, \dots, 0), (1, \dots, 1, 0)). \end{cases}$$

Note that we have $\langle \Theta^\vee, \alpha \rangle = 1$ thus $\langle -\Theta^\vee, \varpi - \alpha \rangle = 0$. We let $v_1 \in V$ be a weight vector of weight ϖ , and $v_d \in V$ of weight $s_\alpha(\varpi) = \varpi - \alpha$. It follows that the line through v_1 and v_d is included in X , and moreover we see that v_1 is the lowest weight of the vector representation of $\tilde{L}_1 \simeq Spin_{\delta+2}$ and v_d is the lowest weight of the representation $\mathbb{S} \otimes W_e$ where $\mathbb{S}$ is a spin representation of $\tilde{L}_1$ and W_e is the natural representation of a simple factor SL_e of the group L_2 (see the following array: the black circles represent the Dynkin diagram, one has to remove the crossed ones to find L_1 ; the blue square represent the longest root and the green disks the weight $\varpi - \alpha$).

X	$\mathbb{P}^{m-1} \times \mathbb{P}^{m-1}$	$\mathbb{G}(2, 2m)$	E_6/P_1
			
d	2	3	5
e	$m - 2$	$2m - 4$	1

It follows that Y'_3 contains the closure S of the $(\tilde{L}_1 \times L_2)$ -orbit of $[v_1 + v_d]$, which is by definition equivalent to a $S(d, e)$. The equality $Y'_3 = S$ follows counting dimensions. In fact, observe from the above array that we always have $e = \frac{d-1}{2}(2m - d) - 1$ (we have no better way to check this than making the computation in the three cases $d = 2, 3, 5$). This may be rewritten as

$$\frac{(d+4)(d-1)}{2} + e - 1 = (d-1)m + 2d - 4. \quad (3)$$

We may find lines in S which meet Y_2 by choosing any point in S and then a line in the $\mathbb{P}^d$ which contains it and was described in Lemma 3.18, taking into account the fact that in this way each line is counted ∞^1 times. The dimension of the set of these lines is thus the left hand-side of (3) since $\dim S = \frac{(d+2)(d-1)}{2} + e$ by Lemma 3.18. On the other hand the variety of lines in X passing through a point in Y_2 has dimension $\dim Y_2 + c_1(X) - 2$, which is the right hand-side of (3) by Lemma 3.15 since $\delta = 2d - 2$. Thus it follows that any line meeting Y_2 is included in S , so that $S = Y'_3$.

Moreover we show that this is also $X \cap \mathbb{P}(T_P \text{Sec}(X))$ by decomposing V as an $(\tilde{L}_1 \times L_2)$ -module (we include a proof for completeness, but this result will not be used in the sequel). Note that we have a decomposition $V = V_1 \oplus V_{1/2} \oplus V_0$ with $V_1, V_{1/2}, V_0$ as in the following array:

δ	2	4	8
V	$A \otimes B$ ($\dim A = \dim B = m$)	$\wedge^2 A$ ($\dim A = 2m$)	V_{27}
with	$A = A_2 \oplus A_{2m-2}, B = B_2 \oplus B_{2m-2}$	$A = A_4 \oplus A_{2m-4}$	
$\tilde{L}_1$	$SL(A_2) \times SL(B_2)$	$SL(A_4)$	$Spin_{10}$
L_2	$SL(A_{m-2}) \times SL(B_{m-2})$	$SL(A_{2m-4})$	$\{1\}$
V_1	$A_2 \otimes B_2$	$\wedge^2 A_4$	V_{10}
$V_{1/2}$	$A_2 \otimes B_{2m-2} \oplus B_2 \otimes A_{2m-2}$	$A_4 \otimes A_{2m-4}$	S_{16}
V_0	$A_{m-2} \otimes B_{m-2}$	$\wedge^2 A_{2m-4}$	$\mathbb{C}$

Assuming P is an idempotent in the Jordan algebra V (which we can always do up to an action of G), the above decomposition is the Pierce decomposition $V = V_1 \oplus V_{1/2} \oplus V_0$, where V_x is the eigenspace for multiplication by P with eigenvalue x .

Note that the linear span of Y_2 is V_1 and that $T_P \text{Sec}(x) = V_1 \oplus V_{1/2}$. Let $y \in Y'_3$: by definition this means that there exists $x \in Y_2$ such that the line (xy) is included in X . Therefore $y \in T_x X$. By Terracini lemma we have $Y_x X \subset T_P \text{Sec}(X)$, so that we deduce $Y'_3 \subset T_P \text{Sec}(X)$. In the case $\delta = 2$ resp. $\delta = 4$, Y'_3 is clearly the set of couples $(x, y) \in \mathbb{P}A \times \mathbb{P}B$ with $x \in \mathbb{P}A_2$ or $y \in \mathbb{P}B_2$ resp. the set of subspaces V_2 meeting non trivially A_4 . Thus this is indeed $X \cap \mathbb{P}(T_P \text{Sec}(X))$. In the case of E_6/P_1 , Y'_3 has dimension 15 and must therefore equal the hyperplane section $\mathbb{P}(T_P \text{Sec}(X)) \cap X$. $\square$

3.6 Y'_3 for a hyperplane section of a Scorza variety

Let X be a Scorza variety and W a general hyperplane section of X . We will reduce the computation of $Y_3(W)$ to the case when $X = \mathbb{P}^1 \times \mathbb{P}^2$ (which is not a Scorza variety), and therefore we start studying this case.

Lemma 3.21 *Let $X = \mathbb{P}^1 \times \mathbb{P}^2$ and W a generic hyperplane section of X . Given 3 points x_1, x_2, x_3 in W and a reducible curve R of degree 3 through these points, R is a limit of some irreducible curves passing through x_1, x_2, x_3 and contained in W .*

Thus the same result holds for a generic curve in X .

Proof. First of all, let us describe W as a scroll. Restricting the linear form H on the $\mathbb{P}^1$'s of $\mathbb{P}^1 \times \mathbb{P}^2$ yields a linear map $\mathbb{C}^3 \rightarrow \mathbb{C}^2$, with 1-dimensional kernel. Let $x \in \mathbb{P}^2$ correspond to this kernel: it is the only element in $\mathbb{P}^2$ such that $\mathbb{P}^1 \times \{x\}$ is contained in H . Take a line l in $\mathbb{P}^2$ not containing x and let $C := (\mathbb{P}^1 \times l) \cap H$. Then C is a conic which may be described as the set of elements $(a, f(a))$ for some isomorphism $f : \mathbb{P}^1 \rightarrow l$. Then W is the union of all the lines joining $(a, f(a)) \in C$ and $(a, x) \in \mathbb{P}^1 \times \{x\}$.

Now, given three generic points x_1, x_2, x_3 on W , we may choose C such that $x_1, x_2 \in C$, so that $x_i = (a_i, f(a_i)) \in \mathbb{P}^1 \times l$ for some $a_i \in \mathbb{P}^1$ and $i = 1, 2$. Then the third point is on the line joining a point $(a_3, f(a_3))$ in C and $(a_3, x) \in \mathbb{P}^1 \times \{x\}$. Thus we may assume that $R = C \cup l$, where l is the line between $(a_3, f(a_3))$ and (a_3, x) .

If h resp. q is a linear resp. quadratic form on $\mathbb{P}^1$ then the morphism

$$t \mapsto (t, h(t)f(t) + q(t)x) \in W \subset \mathbb{P}^1 \times \mathbb{P}^2$$

will pass through x_1, x_2 provided that q vanishes at a_1, a_2 , and imposing moreover that it passes through x_3 yields one more relation between h and q . Thus there are indeed infinitely many irreducible curves of degree 3 through x_1, x_2, x_3 .

If q tends to 0 with the vanishing point of h tending accordingly to a_3 , we see that the limit of those curves must contain C . Since it is connected and contains x_3 , this limit can only be R . $\square$

Let X be a Scorza variety and $W = X \cap H$ a generic hyperplane section of X .

Proposition 3.22 *We have $Y_3(X) = Y'_3(X)$ and $Y_3(W) = Y_3(X) \cap H = Y'_3(W)$, and if there passes a curve of degree 3 through 3 points of W , then there passes infinitely many.*

Proof. We show that any reducible curve R in X or W of degree 3 through x_1, x_2, x_3 is a limit of some irreducible curves passing through the same points, which implies the proposition.

We first consider the case of X . To see this, we use the description of $Y'_3(X)$ given in Proposition 3.20. By an action of G , we may assume that x_1 resp. x_2 is the projectivisation of the highest resp. lowest weight line in V , so that $x_1 = a_1, x_2 = a_2$ are some points on the quadric $Q(d)$, and x_3 is on the line joining a point a_3 in $Q(d)$ and a point $b_3 = s_3 \otimes w_3$ in $S \times \mathbb{P}W$, with $a_3 \in L_d(s_3)$, and finally that R is the union of this line and a conic C included in $Q(d)$ and passing through a_1 and a_2 . Let $V_{d-2} = L_d(s_3) \cap \langle a_1, a_2, a_3 \rangle^\perp$, an isotropic subspace of dimension $d-2$. We choose a non-degenerate subspace V_4 of dimension 4 included in $V_{d-2}^\perp$, containing a_1, a_2, a_3 , and disjoint from V_{d-2} .

Since $\mathbb{P}V_4 \cap Q(d)$ is a smooth quadric surface, to each point $a \in \mathbb{P}V_4 \cap Q(d)$ corresponds an isotropic plane that we denote V_a , such that $\tilde{V}_a := V_{d-2} \oplus V_a$ is a maximal isotropic subspace in V_{2d} parametrised by a point s_a in S . Remark that any two distinct subspaces $\tilde{V}_a, \tilde{V}_b$ meet in codimension 2 (namely, along V_{d-2}), so that the curve traced out by the s_a 's in the spinor variety is in fact a line.

It follows that X contains a $\mathbb{P}^1 \times \mathbb{P}^2$, namely the union of all the planes $\mathbb{P}(V_a \oplus \mathbb{C}.s_a \otimes w_3)$ for a in the quadric surface. Moreover, since $\tilde{V}_{a_3} = L_d(s_3)$, this $\mathbb{P}^1 \times \mathbb{P}^2$ contains the line joining a_3 and b_3 . Since $\mathbb{P}V_4 \cap Q(d)$ contains a_1, a_2, a_3 , it contains C and thus our $\mathbb{P}^1 \times \mathbb{P}^2$ contains our initial curve R . Thus by Lemma 3.21, R can be deformed to an irreducible curve passing through x_1, x_2, x_3 .

If we assume from the beginning that R is included in $W = X \cap H$, then again from Lemma 3.21, the same conclusion holds. $\square$

Theorem 3 *Let X be an adjoint or a coadjoint variety and let σ_u and σ_v be classical cohomology classes. Then if we write*

$$\sigma_u * \sigma_v = \sum_{d,w} c_{u,v}^w(d) q^d \sigma_w$$

we have $c_{u,v}^w(d) = 0$ unless $d \leq 2$.

Proof. First, by Proposition 3.13 and Proposition 3.22, we know that $c_{u,v}^w(3) = 0$. Lemma 3.1 concludes the proof for $c_{u,v}^w(d)$ with $d > 3$. $\square$

Corollary 3.23 *Let X be an adjoint variety, then in the quantum cohomology $QH^*(X, \mathbb{C})$ we have the formula*

$$\text{pt} * \text{pt} = 2q^2 \ell$$

where pt resp. ℓ denotes the class of a point resp. the class of a line in X .

Proof. By degree counting the first quantum terms are of degree 2 in q . Furthermore by the previous result, there are no higher powers of q . We thus have to prove the equality $I_2(\text{pt}, \text{pt}, h) = 2$ where h is the hyperplane class. As we have seen in Proposition 3.11, there is a unique conic through two general points in X . As this conic meets a hyperplane in two points, the result follows. $\square$

4 Quantum to classical principle for degree one Gromov-Witten invariants

In this section, we explain how to apply by now classical technics initiated by A. Buch, A. Kresch and H. Tamvakis [BuKrTa03] to compute degree one Gromov-Witten invariants. We shall here use the presentation of [ChMaPe08]. Most of the arguments are very similar to those of [ChMaPe08, Section 3] and we will therefore not reproduce them.

Let F be the variety of lines on X . This variety is well described in [LaMa03]. In particular, note that this variety is homogeneous for X adjoint but is not homogeneous for X coadjoint and not adjoint. In the last case, the comparison principle is less useful to compute Gromov-Witten invariants on X than to compute intersections on F .

Let us recall some notation from [ChMaPe08]. Points in F will be denoted by ℓ and we shall denote by L the associated line in X . We have an incidence $I = \{(x, \ell) \in X \times F \mid x \in L\}$ between X and F . We denote by p resp. q the projection from I to X resp. F . For $w \in W^P$, we denote the variety $q(p^{-1}(X(w)))$ by $F(\hat{w})$.

The following lemma is an adaptation of Lemma 3.7 in [ChMaPe08] to the non minuscule case. For $w \in W^P$, we denote by $w^\vee$ the element of W^P such that $X(w)$ and $X(w^\vee)$ are Poincaré dual.

Lemma 4.1 *Let $X(w)$ and $X(v^\vee)$ be two Schubert varieties such that $X(w) \subset X(v^\vee)$. Then there exists an element $g \in G$ such that the intersection $X(w) \cap g \cdot X(v)$ is a reduced point.*

Proof. There always exists a sequence of inclusions of Schubert varieties

$$X(w) = X(w_0) \subset X(w_1) \subset \cdots \subset X(w_r) = X(v^\vee)$$

where $X(w_i)$ is a moving divisor in $X(w_{i+1})$ for all i . We prove, by descending induction on $i \in [0, r]$, that there exists an element g_i in G such that the intersection $X(w_i) \cap g_i \cdot X(v)$ is a reduced point. For $i = r$ this is Poincaré duality. Assume that the result holds for w_{i+1} i.e. there exists an element $g_{i+1} \in G$ such that $X(w_{i+1}) \cap g_{i+1} \cdot X(v)$ is a reduced point say x . Recall that because $X(w_i)$ is a moving divisor in $X(w_{i+1})$, we have the equality:

$$X(w_{i+1}) = \bigcup_{h \in \text{Stab}(X(w_{i+1}))} h \cdot X(w_i).$$

We deduce from this that there exists an element $h \in \text{Stab}(X(w_{i+1}))$ such that $x \in h \cdot X(w_i)$. But now $h \cdot X(w_i)$ meets $g_{i+1} \cdot X(v)$ in x at least. Since $h \cdot X(w_i)$ is contained in $X(w_{i+1})$ and $X(w_{i+1}) \cap g_{i+1} \cdot X(v)$ is the reduced point x , we have that $h \cdot X(w_i)$ and $g_{i+1} \cdot X(v)$ meet transversely and only in x . $\square$

For the next two results, the proofs of [ChMaPe08, Lemma 3.10 and Corollary 3.11] apply verbatim.

Lemma 4.2 *Let $X(u)$, $X(v)$ and $X(w)$ be three proper Schubert subvarieties of X , such that*

$$\text{Codim}(X(u)) + \text{Codim}(X(v)) + \text{Codim}(X(w)) = \dim(X) + c_1(X).$$

Then for g, g' and g'' three general elements in G , the intersection $g \cdot F(\hat{u}) \cap g' \cdot F(\hat{v}) \cap g'' \cdot F(\hat{w})$ is a finite set of reduced points.

Let ℓ be a point in this intersection, then the line L meets each of $g \cdot X(u)$, $g' \cdot X(v)$ and $g'' \cdot X(w)$ in a unique point and these points are in general position in L .

Corollary 4.3 *Let $X(u)$, $X(v)$ and $X(w)$ be three proper Schubert subvarieties of X . Suppose that the sum of their codimensions is $\dim(X) + c_1(X)$. Then*

$$I_1([X(u)], [X(v)], [X(w)]) = I_0([F(\hat{u})], [F(\hat{v})], [F(\hat{w})]).$$

As in [ChMaPe08], we may use this result to compute combinatorially some Gromov-Witten invariants. Let us simply give an example of this.

Example 4.4 We assume that X is of adjoint type and homogeneous under a group of type different from A_n . Let u and v be elements in W^P such that $l(u) + l(v) = c_1(X) = \deg(q)$. We want to compute the quantum product $\sigma^u * \sigma^v$.

Let us first remark that in this situation, if $c_1 = c_1(X)$, then we have $\dim X = 2c_1 - 1$. In particular, the only quantum non classical term appearing in $\sigma^u * \sigma^v$ is of the form aq for some $a \in \mathbb{Z}$. In symbols:

$$\sigma^u * \sigma^v = \sigma^u \cup \sigma^v + aq.$$

The integer a is equal to the Gromov-Witten invariant $I_1([X(u^\vee)], [X(v^\vee)], [\text{pt}])$. By the previous discussion, this invariant is equal to $I_0([q(p^{-1}(X(u^\vee)))], [q(p^{-1}(X(v^\vee)))], [q(p^{-1}(\text{pt}))])$.

In our situation, the variety F is homogeneous of the form G/Q for Q a parabolic subgroup. By the same discussion as in [ChMaPe08, Subsection 3.3], the variety $F(\hat{w}) = q(p^{-1}(X(w)))$ for $X(w)$ a Schubert variety in X is a Schubert variety in F . The notation $\hat{w}$ will in that situation stand for the corresponding element in W^Q . Furthermore, the incidence variety I is also homogeneous of the form G/R with $R = P \cap Q$ and the inverse image $p^{-1}(X(w))$ of a Schubert variety is again a Schubert variety. We may obtain the element $\hat{w}$ easily from w by the following procedure.

- First remark that P is maximal associated to a simple root, say α , and that w_X the longest element in W^P has the form $s_\alpha \tilde{w}_X$ with $l(\tilde{w}_X) = l(w_X) - 1$. In other words $\tilde{w}_X = s_\alpha w_X$.
- Let Z denote the fiber $p^{-1}(\text{pt})$ of the zero dimensional Schubert variety pt in X . This is a Schubert variety in I and let $w_Z \in W^R$ be the corresponding element. Then $p^{-1}(X(w))$ is the Schubert variety in I associated to $ww_Z \in W^R$.
- If $w \neq w_X$, then we have $\hat{w} = ww_Z \in W^Q$. We have $\hat{w}_X = s_\alpha w_X w_Z$. This last element is the longest element in W^R , we denote it by w_F . This element is self-inverse thus $w_F = w_Z w_X s_\alpha$.

We want to compute the Littlewood-Richardson coefficient

$$I_0([F(\widehat{u^\vee})], [F(\widehat{v^\vee})], [F(w_Z)]) = c_{\widehat{u^\vee}^\star, \widehat{v^\vee}^\star}^{w_Z}$$

in F with $\star$ denoting Poincaré duality in F . Now compute for $l(u) > 0$:


$$\widehat{u^\vee}^\star = w_0 \widehat{u^\vee} w_0 w_F = w_0 (u^\vee w_Z) w_0 w_F = w_0 (w_0 u w_0 w_X) w_Z w_0 w_F = u w_X w_Z w_F = u s_\alpha.$$

The same is true for $\widehat{v^\vee}^\star$. We thus get

$$a = c_{u s_\alpha, v s_\alpha}^{w_Z}.$$

Now remark that for X adjoint the fiber Z is always a product of minuscule and cominuscule homogeneous spaces. In particular of Λ is the weight associated to Q , then w_Z is Λ -minuscule and we may compute the above Littlewood-Richardson coefficient using jeu de taquin by [ChPe09a, Theorem 1.3].

Let us do an explicit example. Let $X = E_6/P_2$, the heap associated to w_Z is the heap of a Grassmannian as follows:



Let $u = s_{\alpha_2} s_{\alpha_4} s_{\alpha_5} s_{\alpha_3} s_{\alpha_4} s_{\alpha_2} \in W^{P_2}$ and v be any element in W^{P_2} with $l(v) = 5$. We have $\alpha = \alpha_2$ and $u s_\alpha = s_{\alpha_2} s_{\alpha_4} s_{\alpha_5} s_{\alpha_3} s_{\alpha_4}$, thus its heap is not contained in the heap of w_Z and we get

$$\sigma^u * \sigma^v = \sigma^u \cup \sigma^v.$$

All the coefficients in this classical product can be computed using jeu de taquin.

Let us now set $u' = s_{\alpha_1} s_{\alpha_4} s_{\alpha_5} s_{\alpha_3} s_{\alpha_4} s_{\alpha_2} \in W^{P_2}$ and $v' s_{\alpha_1} s_{\alpha_5} s_{\alpha_3} s_{\alpha_4} s_{\alpha_2} \in W^{P_2}$ be elements in W^{P_2} . We have $u' s_\alpha = s_{\alpha_1} s_{\alpha_4} s_{\alpha_5} s_{\alpha_3} s_{\alpha_4}$ and $v' s_\alpha = s_{\alpha_1} s_{\alpha_5} s_{\alpha_3} s_{\alpha_4}$, thus by jeu de taquin we get

$$\sigma^u * \sigma^v = \sigma^u \cup \sigma^v + q.$$

Again, all the coefficients in the classical product $\sigma^u \cup \sigma^v$ can be computed using jeu de taquin.

5 Presentation of the quantum cohomology rings

Recall that for the minuscule and cominuscule varieties, presentation of the quantum cohomology rings are already known: in type A this is done in [SiTi97], in type B , C and D in [KrTa03] and [KrTa04]. In [ChMaPe08] the authors gave a uniform presentation of the quantum cohomology algebras of the minuscule and cominuscule homogeneous spaces. In this section we give a presentation of the quantum cohomology algebras of adjoint and coadjoint homogeneous spaces. One more time for classical groups this is already known: in type A , a presentation will follow from the quantum Chevalley formula (see Section 7), in type B , C and D , such a presentation is contained in the more general results of [BuKrTa08]. Therefore in this section we are concerned with exceptional types.

Most of the computations here made use of a software written in Java, that the reader may upload or test at the webpage www.math.sciences.univ-nantes.fr/~chaput/quiver-demo.html, in order to visualise the quantum Hasse diagrams, apply our version of the Littlewood-Richardson rule [ChPe09a], perform quantum multiplications, and in general check the assertions in this section.

In the proofs in this section, the Hasse diagrams are colored: the orange vertices correspond to ϖ -minuscule elements. The blue vertices correspond to quantum Schubert classes with a factor q while grey vertices correspond to quantum Schubert classes with a factor q^2 . In the paper [ChPe09b], we gathered the quantum Giambelli formulas for all exceptional adjoint and coadjoint varieties. We also give in [ChPe09b] some tableaux used in the proof of the presentation for E_8/P_8 (we did not include these tableaux in this paper to save place).


5.1 Adjoint and coadjoint varieties in type G_2

The homogeneous space G_2/P_1 is a smooth 5-dimensional quadric, thus its quantum cohomology ring is $QH^*(G_2/P_1, \mathbb{Q}) = \mathbb{Q}[h, q]/(4hq - h^6)$. For G_2/P_2 we have the following:

Proposition 5.1 *Let h denote the hyperplane class of G_2/P_2 . We have*

$$QH^*(G_2/P_2) = \mathbb{Q}[h, q]/(h^6 - 18h^3q - 27q^2).$$

Proof. Using Theorem 2, it is immediate to check that the quantum Hasse diagram of G_2/P_2 begins as follows (and then it is periodic):




From this one computes that the degree 2 class is $\sigma_2 = 1/3h^2$ and that the class of a point is $\sigma_5 = 1/18h^5 - 5/6h^2q$. We deduce from this that $q^2 = (2h\sigma_5 - hq\sigma_2)/3 = 1/27h^6 - 2/3h^3q$. $\square$

5.2 Adjoint and coadjoint varieties in type F_4

Proposition 5.2 *Let h denote the hyperplane class of F_4/P_1 and $s \in H^*(F_4/P_1)$ be the Schubert class corresponding to $w = s_2 s_3 s_2 s_1$. We have*

$$QH^*(F_4/P_1, \mathbb{Q}) = \mathbb{Q}[h, s, q]/(h^8 - 12s^2 - 16q, 3h^{12} - 18h^8s + 24h^4s^2 + 8s^3).$$

Proof. The quantum Hasse diagram of F_4/P_1 begins as follows:




To compute s^2 in $QH^*(F_4/P_1)$, we first compute the classical product. Let $\sigma_{4,2}$ correspond to $s_4s_3s_2s_1$: the Hasse diagram shows that $\sigma_{4,2}^2$ has degree 56. Let $\sigma_{8,1}$ resp. $\sigma_{8,2}$ correspond to $s_1s_2s_3s_2s_4s_3s_2s_1$ resp. $s_2s_1s_3s_2s_4s_3s_2s_1$. Since $\deg(\sigma_{8,1}) = 16$ and $\deg(\sigma_{8,2}) = 40$, we deduce that $\sigma_{4,2} \cup \sigma_{4,2} = \sigma_{8,1} + \sigma_{8,2}$. Using the Chevalley formula it follows that $s \cup s = 6\sigma_{8,1} + 8\sigma_{8,2}$. The quantum product $s * s = s^2$ is then equal to the same class by Example 4.4. By quantum Chevalley formula, we get the first relation. The second relation (of degree 12) is then just a consequence of the Chevalley formula. This is enough in order to express all the Schubert classes as polynomials in h, s and q , as displayed in [ChPe09b]. $\square$

Proposition 5.3 *Let h denote the hyperplane class of F_4/P_4 and $s \in H^*(F_4/P_1)$ be the Schubert class corresponding to $w = s_1s_2s_3s_4$. We have*

$$QH^*(F_4/P_4, \mathbb{Q}) = \mathbb{Q}[h, s, q] / (2h^8 - 6h^4s + 3s^2, -11h^{12} + 26h^8s + 3hq).$$

Proof. The quantum Hasse diagram of F_4/P_4 begins as follows:




This Hasse diagram shows that the classical (or quantum because its cohomological degree is $8 < c_1$) product $s \cup s = s * s$ has degree 14. Let us denote $\sigma_{8,1}$ resp. $\sigma_{8,2}$ the Schubert classes corresponding to $s_3s_2s_4s_1s_3s_2s_3s_4$ resp. $s_4s_3s_2s_1s_3s_2s_3s_4$, of degree 5 resp. 2. We have either $s^2 = 2\sigma_{8,1} + 2\sigma_{8,2}$ or $s^2 = 7\sigma_{8,2}$. Since $s * h^4 \geq s^2$ and $s * h^4 = 5\sigma_{8,1} + 4\sigma_{8,2}$, the second case is excluded. Thus we have $s^2 = 2\sigma_{8,1} + 2\sigma_{8,2}$. The relations in degree 8 and 12 follow. $\square$

5.3 Adjoint and coadjoint varieties in type E

We now consider the technically more involved cases of groups of type E . In the cohomology of E_6/P_2 , let h denote the hyperplane class, s resp. t the classes corresponding to the elements $s_3s_4s_2$ resp. $s_1s_3s_4s_2$.

Proposition 5.4 *The quantum cohomology algebra $QH^*(E_6/P_2, \mathbb{Q})$ is the quotient of the polynomial algebra $\mathbb{Q}[h, s, t, q]$ by the relations $h^8 - 6h^5s + 3h^4t + 9h^2s^2 - 12hst + 6t^2$, $h^9 - 4h^6s + 3h^5t + 3h^3s^2 - 6h^2st + 2s^3$, and $-97h^{12} + 442h^9s - 247h^8t - 507h^6s^2 + 624h^5st - 156h^2s^2t + 48hq$.*

Proof. We know that $QH^*(E_6/P_2, \mathbb{Q})$ is generated by h, s and t and that there are relations in degree 8, 9, 12. The quantum Hasse diagram of E_6/P_2 begins as follows:




From this it follows that the degree of s^2 is 37752. From now on and until the end of this section, we will write Schubert classes according to the following notation in the proofs:

Notation 5.5 To $w \in W/W_P$ we associate the element $\varphi(w) = \alpha = \varpi - w(\varpi)$ of the root lattice. If $\alpha = \varphi(w)$ with $w \in W/W_P$, we denote $\sigma(\alpha)$ the cohomology class σ_w .

Let us define the following Schubert classes

$$\sigma_{6,1} = \sigma\left(\begin{smallmatrix} 11210 \\ 2 \end{smallmatrix}\right), \sigma_{6,2} = \sigma\left(\begin{smallmatrix} 01210 \\ 2 \end{smallmatrix}\right), \sigma_{6,3} = \sigma\left(\begin{smallmatrix} 11111 \\ 1 \end{smallmatrix}\right), \text{ and } \sigma_{6,4} = \sigma\left(\begin{smallmatrix} 01211 \\ 1 \end{smallmatrix}\right),$$

they have respective degrees 10920, 6006, 4992, and 10920. Considering the following tableaux we deduce from [ChPe09a] that $s^2 \geq 2\sigma_{6,1} + \sigma_{6,3} + \sigma_{6,4}$:




On the other hand both s^2 and $2\sigma_{6,1} + \sigma_{6,3} + \sigma_{6,4}$ have degree 37752: thus we have $s^2 = 2\sigma_{6,1} + \sigma_{6,3} + \sigma_{6,4}$. This allows to compute an expression of all Schubert classes up to degree 6 as polynomials in h, s, t , as displayed in [ChPe09b].

Let $\sigma_{7,i}, 1 \leq i \leq 5$ be the Schubert classes in $H^7(E_6/P_2)$, corresponding respectively to the roots

$$\left(\begin{smallmatrix} 11210 \\ 2 \end{smallmatrix}\right), \left(\begin{smallmatrix} 12210 \\ 1 \end{smallmatrix}\right), \left(\begin{smallmatrix} 11211 \\ 1 \end{smallmatrix}\right), \left(\begin{smallmatrix} 01211 \\ 2 \end{smallmatrix}\right), \text{ and } \left(\begin{smallmatrix} 01221 \\ 1 \end{smallmatrix}\right).$$


They have respective degrees 3003, 2925, 4992, 3003, and 2925. The product st has degree 7917. In view of the following two tableaux it follows that $st = \sigma_{7,2} + \sigma_{7,3}$.



This is enough to express some Giambelli formulas up to degree 9 but to get the relation in degree 8 resp. 9 we must compute t^2 resp. s^3 . To this end we compute t^2 and $s \cdot \sigma_{6,1}$. The degree of t^2 and

$$\sigma\left(\begin{smallmatrix} 12211 \\ 1 \end{smallmatrix}\right)$$

are 1638, and we have the following tableau:




thus

$$t^2 = \sigma\left(\begin{smallmatrix} 12211 \\ 1 \end{smallmatrix}\right).$$

The following seven tableaux prove the inequality


$$s \cdot \sigma_{6,1} \geq \sigma\left(\begin{smallmatrix} 12310 \\ 2 \end{smallmatrix}\right) + 3\sigma\left(\begin{smallmatrix} 12211 \\ 2 \end{smallmatrix}\right) + \sigma\left(\begin{smallmatrix} 11221 \\ 1 \end{smallmatrix}\right) + 2\sigma\left(\begin{smallmatrix} 12221 \\ 1 \end{smallmatrix}\right) :$$



We leave it to the reader to check that the degrees coincide, so that we have equality.

To express the Schubert classes in $H^{10}(E_6/P_2)$ as polynomials in h, s, t , we must compute s^2t , and to this end we compute the product $t\sigma_{6,1}$. According to the following three tableaux and using the same degree argument as above this product is

$$\sigma\left(\begin{smallmatrix} 12311 \\ 2 \end{smallmatrix}\right) + \sigma\left(\begin{smallmatrix} 12221 \\ 2 \end{smallmatrix}\right) + \left(\begin{smallmatrix} 12321 \\ 1 \end{smallmatrix}\right).$$




These computations are enough to express all the Schubert classes as polynomials in h, s, t, q , and the displayed relation in degree 12 comes for free. $\square$

5.4 Adjoint and coadjoint varieties in type E_7

In the cohomology of E_7/P_1 , let h denote the hyperplane class, s resp. t the classes corresponding to the elements $s_2s_4s_3s_1$ resp. $s_7s_6s_5s_4s_3s_1$.

Proposition 5.6 *The quantum cohomology algebra $QH^*(E_7/P_1, \mathbb{Q})$ is the quotient of the polynomial algebra $\mathbb{Q}[h, s, t, q]$ by the relations $h^{12} - 6h^8s - 4h^6t + 9h^4s^2 + 12h^2st - s^3 + 3t^2$, $h^{14} - 6h^{10}s - 2h^8t + 9h^6s^2 + 6h^4st - h^2s^3 + 3s^2t$, and $232h^{18} - 1444h^{14}s - 456h^{12}t + 2508h^{10}s^2 + 1520h^8st - 988h^6s^3 + 133h^2s^4 + 36hq$.*


Proof. We know that $QH^*(E_7/P_1, \mathbb{Q})$ is generated by h, s and t and that there are relations in degree 12, 14, 18. The quantum Hasse diagram of E_7/P_1 begins as follows:




The strategy of proof is the same as in the previous cases; therefore we will only sketch the arguments. We can express the Giambelli formulas up to degree 7 without computing any product. In degree 8 we need to compute s^2 . We have

$$s^2 = \sigma\left(\begin{smallmatrix} 222100 \\ 1 \end{smallmatrix}\right) + \sigma\left(\begin{smallmatrix} 122110 \\ 1 \end{smallmatrix}\right) + \sigma\left(\begin{smallmatrix} 112210 \\ 1 \end{smallmatrix}\right) + \sigma\left(\begin{smallmatrix} 112111 \\ 1 \end{smallmatrix}\right).$$

In fact, computing degrees, this follows from the existence of the following four tableaux:



In degree 10, we have to compute st : this is equal to $\sigma\left(\begin{smallmatrix} 122211 \\ 1 \end{smallmatrix}\right)$, in view of:




In degree 12, we have two new monomials: t^2 and s^3 . We have

$$t^2 = \sigma\left(\begin{smallmatrix} 222221 \\ 1 \end{smallmatrix}\right).$$

To express s^3 as a sum of Schubert classes and thus find the given relation of degree 12, we compute that

$$s \cdot \sigma\left(\begin{smallmatrix} 222100 \\ 1 \end{smallmatrix}\right) = \sigma\left(\begin{smallmatrix} 233210 \\ 1 \end{smallmatrix}\right) + \sigma\left(\begin{smallmatrix} 223211 \\ 1 \end{smallmatrix}\right) + \sigma\left(\begin{smallmatrix} 222221 \\ 1 \end{smallmatrix}\right)$$

The corresponding tableau for t^2 is displayed on the left and the three tableaux for the above product are on the right:




To get the relation in degree 14, we have to compute the monomial s^2t . To this end we compute that

$$t \cdot \sigma \begin{pmatrix} 222100 \\ 1 \end{pmatrix} = \sigma \begin{pmatrix} 223221 \\ 2 \end{pmatrix}.$$

To express the Schubert classes in $H^{16}(E_7/P_1)$, we need to compute st^2 , and to this end we show that

$$t \cdot \sigma \begin{pmatrix} 112221 \\ 1 \end{pmatrix} = \sigma \begin{pmatrix} 233321 \\ 2 \end{pmatrix}.$$

These relations come from the two tableaux:



This is enough to express all Schubert classes as polynomials in h, s, t, q , and yields the last relation.


□

5.5 Adjoint and coadjoint varieties in type E_8

In the cohomology of E_8/P_8 , let h denote the hyperplane class, s resp. t the classes corresponding to the elements $s_2s_4s_5s_6s_7s_8$ resp. $s_6s_5s_4s_3s_2s_4s_5s_6s_7s_8$.

Proposition 5.7 *The quantum cohomology algebra $QH^*(E_8/P_8, \mathbb{Q})$ is the quotient of the polynomial algebra $\mathbb{Q}[h, s, t, q]$ by the relations $h^{14}s + 6h^{10}t - 3h^8s^2 - 12h^4st - 10h^2s^3 + 3t^2$, $29h^{24} - 120h^{18}s + 15h^{14}t + 45h^{12}s^2 - 30h^8st + 180h^6s^3 - 30h^2s^2t + 5s^4$, and $-86357h^{30} + 368652h^{24}s - 44640h^{20}t - 189720h^{18}s^2 + 94860h^{14}st - 473680h^{12}s^3 + 74400h^8s^2t - 1240h^2s^3t + 60hq$.*

Proof. We know that $QH^*(E_8/P_8, \mathbb{Q})$ is generated by h, s and t and that there are relations in degree 14, 24, 30. The quantum Hasse diagram of E_8/P_8 begins as follows:



From this we may compute the degrees of all the Schubert classes and of any product of two Schubert classes. Moreover we argue as before to compute such products, and display the results on the companion paper [ChPe09b]. We only indicate the computed products, and why they suffice to obtain the presentation. Note that these results were obtained with the help of a computer by S. Nikolenko and N. Semenov [NiSe06].

We have

$$s^2 = 2\sigma\left(\begin{smallmatrix} 1232111 \\ 1 \end{smallmatrix}\right) + 4\sigma\left(\begin{smallmatrix} 1222211 \\ 1 \end{smallmatrix}\right) + 2\sigma\left(\begin{smallmatrix} 1122221 \\ 1 \end{smallmatrix}\right) + \sigma\left(\begin{smallmatrix} 0122222 \\ 1 \end{smallmatrix}\right) \text{ and}$$

$$s \cdot t = \sigma\left(\begin{smallmatrix} 1243211 \\ 2 \end{smallmatrix}\right) + 5\sigma\left(\begin{smallmatrix} 1233221 \\ 2 \end{smallmatrix}\right) + 6\sigma\left(\begin{smallmatrix} 1232222 \\ 2 \end{smallmatrix}\right) + 2\sigma\left(\begin{smallmatrix} 1233321 \\ 1 \end{smallmatrix}\right) + 7\sigma\left(\begin{smallmatrix} 1233222 \\ 1 \end{smallmatrix}\right).$$

This allows to compute some Giambelli formulas up to degree 17. To compute s^3 we compute

$$s \cdot \sigma\left(\begin{smallmatrix} 0122222 \\ 1 \end{smallmatrix}\right) = \sigma\left(\begin{smallmatrix} 1243222 \\ 2 \end{smallmatrix}\right) + \sigma\left(\begin{smallmatrix} 1233322 \\ 2 \end{smallmatrix}\right) + \sigma\left(\begin{smallmatrix} 1233332 \\ 1 \end{smallmatrix}\right),$$

which allows to get the Giambelli formulas up to degree 21. To get the relation in degree 20 we compute

$$t^2 = 4\sigma\left(\begin{smallmatrix} 2343321 \\ 2 \end{smallmatrix}\right) + 2\sigma\left(\begin{smallmatrix} 1344321 \\ 2 \end{smallmatrix}\right) + 7\sigma\left(\begin{smallmatrix} 2343222 \\ 2 \end{smallmatrix}\right) + 14\sigma\left(\begin{smallmatrix} 1343322 \\ 2 \end{smallmatrix}\right) \\ + 8\sigma\left(\begin{smallmatrix} 1244322 \\ 2 \end{smallmatrix}\right) + 16\sigma\left(\begin{smallmatrix} 1243332 \\ 2 \end{smallmatrix}\right).$$

To express the Schubert classes of degree 22 we need to compute s^2t : to this end we compute

$$t \cdot \sigma\left(\begin{smallmatrix} 0122222 \\ 1 \end{smallmatrix}\right) = \sigma\left(\begin{smallmatrix} 2344322 \\ 2 \end{smallmatrix}\right) + 3\sigma\left(\begin{smallmatrix} 1354332 \\ 3 \end{smallmatrix}\right) + 2\sigma\left(\begin{smallmatrix} 2344432 \\ 2 \end{smallmatrix}\right) + \sigma\left(\begin{smallmatrix} 1243332 \\ 2 \end{smallmatrix}\right).$$

To get the relation in degree 24 we need to know s^4 , and for this we show

$$\sigma\left(\begin{smallmatrix} 0122222 \\ 1 \end{smallmatrix}\right)^2 = \sigma\left(\begin{smallmatrix} 2344432 \\ 2 \end{smallmatrix}\right).$$

Finally we need to express the Schubert classes of degree 28 in terms of the generators; for this we need to compute s^3t which follows from the equality

$$\sigma\left(\begin{smallmatrix} 0122222 \\ 1 \end{smallmatrix}\right) \cdot \sigma\left(\begin{smallmatrix} 1233222 \\ 1 \end{smallmatrix}\right) = \sigma\left(\begin{smallmatrix} 2464432 \\ 3 \end{smallmatrix}\right) + \sigma\left(\begin{smallmatrix} 2465432 \\ 2 \end{smallmatrix}\right) + 4\sigma\left(\begin{smallmatrix} 2455432 \\ 3 \end{smallmatrix}\right) + \sigma\left(\begin{smallmatrix} 2365432 \\ 3 \end{smallmatrix}\right).$$

The relation in degree 30 follows from the Chevalley formula. $\square$

6 Semi-simplicity of the quantum cohomology and strange duality

Recall that in [ChMaPe07] and [ChMaPe09a] we proved that the localisation

$$QH^*(X, \mathbb{C})_{loc} := QH^*(X, \mathbb{C})[q^{-1}]$$

of the quantum parameter in the quantum cohomology algebra of any minuscule or cominuscule variety X is semi-simple. A general argument using complex conjugation (see Theorem 2.1 in [ChMaPe09a]), implies that in that case there exists an algebra involution on $QH^*(X, \mathbb{C})_{loc}$ sending q on its inverse and more generally a quantum cohomology class of degree d to a quantum cohomology class of degree $-d$. We conjectured in [ChMaPe09a] that the only rational homogeneous spaces X with Picard number one such that $QH^*(X, \mathbb{C})_{loc}$ is semi-simple were the minuscule and cominuscule ones.

6.1 Coadjoint varieties

Theorem 4 *Assume that there exists a presentation of the classical cohomology algebra $H^*(X, \mathbb{C})$ satisfying:*

- *there is no equation of degree multiple of $c_1(X)$,*
- *there is at least one equation of degree smaller than $c_1(X)$.*

Then the localised quantum cohomology algebra $QH^(X, \mathbb{C})$ is not semi-simple.*

Proof. First remark that to prove that the localised quantum cohomology algebra $QH^*(X, \mathbb{C})$ is not semi-simple, it suffices to prove that the specialisation of the quantum cohomology algebra at $q = 1$ is not semi-simple.

Let us fix some notation for the presentation of the cohomology ring $H^*(X, \mathbb{C})$. This ring can always be presented as a homogeneous complete intersection ring: the generators and the equations are homogeneous and there are as many equations as there are generators. Let us denote by $(X_i)_{i \in [1, s]}$ the generators and with $(F_i)_{i \in [1, s]}$ the equations. The generators will be called classical generators and the equations classical equations.

A presentation for the quantum cohomology algebra can be derived from a presentation of the classical cohomology algebra by taking the same classical generators together with q as generators and by deforming the classical relations F_i in relations G_i of the same degree but eventually with q terms (see [SiTi97]). By hypothesis we get homogeneous equations of degree different from multiples of c_1 . In particular any monomial with a q term is also multiple of a classical generator. This implies that $X_i = 0$ for all $i \in [1, s]$ and $q = 1$ is a solution of the equations $((G_i = 0)_{i \in [1, s]}, q = 1)$. Denote by 0 this point. We want to prove that 0 is a multiple solution of the system $((G_i = 0)_{i \in [1, s]}, q = 1)$.

For this, because the specialisation at $q = 1$ of the quantum algebra $QH^*(X, \mathbb{C})$ is a complete intersection algebra of dimension 0, we only need to prove that the family $(d_0 G_i)_{i \in [1, s]}$ of differentials is not linearly independant. But an equation F_i is not deformed if $\deg(F_i) < \deg(q) = c_1(X)$. In that case $G_i = F_i$ and by minimality F_i is at least quadratic in the variables $(X_i)_{i \in [1, s]}$ therefore its differential $d_0 F_i$ vanishes. In particular the family $(d_0 G_i)_{i \in [1, s]}$ of differentials is not linearly independant and 0 is a multiple solution. $\square$

Corollary 6.1 *Let X be a rational homogeneous space with Picard group $\mathbb{Z}$. Then the localised quantum algebra $QH^*(X, \mathbb{C})_{loc}$ is not semi-simple except maybe in the following cases:*

Type	Variety
A_n	$\mathbb{G}(p, n+1)$ for all $p \in [1, n]$.
B_n	$\mathbb{G}_Q(p, 2n+1)$ for all $p \in [1, n]$ with p even, for $p = 1$ and $p = n$.
C_n	$\mathbb{G}_\omega(p, 2n)$ for all $p \in [1, n]$ with p odd and for $p = n$.
D_n	$\mathbb{G}_Q(p, 2n)$ for all $p \in [1, n-1]$ with p odd and for $p = n$.
E_6	E_6/P_1 and E_6/P_3 (and E_6/P_6 and E_6/P_5 which are isomorphic to the previous ones).
E_7	E_7/P_2 , E_7/P_4 , E_7/P_5 and E_7/P_7 .
E_8	E_8/P_4 and E_8/P_6 .
F_4	F_4/P_1 and F_4/P_2 .
G_2	G_2/P_1 and G_2/P_2 .

Remark 6.2 We already proved in [ChMaPe09a] that for X minuscule or cominuscule the localised quantum cohomology ring $QH^*(X, \mathbb{C})_{loc}$ is semi-simple. In the next subsection we shall prove that the same result holds for adjoint non coadjoint varieties. The only cases left are therefore the following ones (for G_2 the coadjoint variety is a quadric):

Type	Variety
B_n	$\mathbb{G}_Q(p, 2n+1)$ for all $p \in [1, n-1]$ with p even.
C_n	$\mathbb{G}_\omega(p, 2n)$ for all $p \in [2, n-1]$ with p odd.
D_n	$\mathbb{G}_Q(p, 2n)$ for all $p \in [2, n-1]$ with p odd.
E_6	E_6/P_3 (and E_6/P_5 which is isomorphic to the previous one).
E_7	E_7/P_2 , E_7/P_4 and E_7/P_5 .
E_8	E_8/P_4 and E_8/P_6 .
F_4	F_4/P_2 .

Corollary 6.3 *For X a coadjoint variety for a semi-simple group different from G_2 , the localised quantum cohomology algebra $QH^*(X, \mathbb{C})$ is not semi-simple.*

6.2 Adjoint non coadjoint varieties

There are three adjoint non coadjoint varieties: $\mathbb{G}_Q(2, 2n+1)$, F_4/P_1 and G_2/P_2 . In this subsection we prove the following

Theorem 5 *For X an adjoint non coadjoint rational homogeneous space, the localised quantum cohomology algebra $QH^*(X, \mathbb{Q})_{loc}$ is semi-simple. It therefore admits an algebra involution ι sending q to its inverse.*

We prove this result by a case by case analysis. We start with type B_n :

Proposition 6.4 *The localised quantum cohomology algebra $QH^*(\mathbb{G}_Q(2, 2n+1), \mathbb{Q})_{loc}$ is semi-simple. It therefore admits an algebra involution sending q to its inverse.*

Proof. A presentation of the ring $QH^*(\mathbb{G}_Q(2, 2n+1), \mathbb{Z})$ (over the integers) has been given in [BuKrTa08]. We will use a simplified presentation which is only valid over $\mathbb{Q}$ but this is enough for our purpose.

Let us denote by K resp. Q the tautological subbundle resp. quotient bundle on $\mathbb{G}_Q(2, 2n+1)$. The rank of K is 2 while the rank of Q is $2n-1$. Pulling these vector bundles to G/T we get decompositions into line bundles of the pull-backs. We denote x_1, x_2 the Chern classes of the two line bundles corresponding to K . Because the pull-back is injective in cohomology, we may therefore write

$$c_t(K) = (1 + x_1 t)(1 + x_2 t) \quad \text{and} \quad c_t(Q) = (1 - x_1 t)(1 - x_2 t) \prod_{i=3}^n (1 - x_i^2 t^2),$$

where c_t denotes the Chern polynomial. The tautological exact sequence gives $c_t(K)c_t(Q) = 1$. We can compute explicit formulas for the Chern classes of K and Q . Let us write $c_i = c_i(K)$ for $i \in [0, 2]$ and $\psi_j = c_j(Q)$ for $j \in [0, n-1]$. We have the equalities:

$$\begin{aligned} c_0 &= \psi_0 = 1, \quad c_1 = x_1 + x_2, \quad c_2 = x_1 x_2, \\ \psi_{2j} &= (-1)^j \sigma_j(x_3^2, \dots, x_n^2) + (-1)^{j-1} x_1 x_2 \sigma_{j-1}(x_3^2, \dots, x_n^2), \\ \psi_{2j+1} &= (-1)^{j+1} (x_1 + x_2) \sigma_j(x_3^2, \dots, x_n^2), \end{aligned}$$

where σ_j is the j -th elementary symmetric function in $n-2$ variables. Remark that $\sigma_{n-1}(x_3^2, \dots, x_n^2) = 0$ thus we have $\psi_{2n-2} = (-1)^n \sigma_{n-2}(x_3^2, \dots, x_n^2)$ and $\psi_{2n-1} = 0$. This yields a presentation of the cohomology algebra

$$H^*(\mathbb{G}_Q(2, 2n+1), \mathbb{Q}) = \mathbb{Q}[c_1, c_2, (\psi_j)_{j \in [1, 2n-2]}] / (\Sigma_k(x_i^2)_{k \in [1, n]}), \quad (4)$$

where Σ_j is the j -th elementary symmetric function in n variables. This is the Borel presentation, see [Bor53] or [BeGeGe73]. Using the first $n - 2$ relations we can express the Chern classes $(\psi_j)_{j \in [1, 2n-2]}$ in terms of c_1 and c_2 . We get for example the following formulas for $j \in [1, n - 2]$:

$$\sigma_j(x_3^2, \dots, x_n^2) = (-1)^j \sum_{k=0}^j x_1^{2k} x_2^{2(j-k)}.$$

In particular, over $\mathbb{Q}$, only the last two equations are worth defining a presentation of the cohomology ring. These equations are:

$$\psi_{2n-2} + c_1 \psi_{2n-3} + c_2 \psi_{2n-4} = 0 \quad \text{and} \quad c_2 \psi_{2n-2} = 0. \quad (5)$$

In these expressions, the ψ terms can be removed but we shall keep them for the moment as they are easier to express in terms of Schubert varieties. This will be useful to deform these equations in the quantum cohomology ring. Let us describe the classes ψ_{2n-4} , ψ_{2n-3} and ψ_{2n-2} as Schubert varieties. This is classical (see for example [BuKrTa08]). Let $(W_p)_{p \in [1, n]}$ be a complete isotropic flag, we have the equalities:

$$\begin{aligned} \psi_{2n-4} &= 2[\{V_2 \in \mathbb{G}_Q(2, 2n+1) \mid \dim(W_3 \cap V_2) \geq 1\}] \\ \psi_{2n-3} &= 2[\{V_2 \in \mathbb{G}_Q(2, 2n+1) \mid \dim(W_2 \cap V_2) \geq 1\}] \\ \psi_{2n-2} &= 2[\{V_2 \in \mathbb{G}_Q(2, 2n+1) \mid \dim(W_1 \cap V_2) \geq 1\}] \end{aligned}$$

where $[Z]$ denotes the cohomology class of Z . In terms of roots as described in Section 2 (with notation as in [Bou54]), we have $\psi_{2n-4} = 2\sigma_{\alpha_1 + \alpha_2}$, $\psi_{2n-3} = 2\sigma_{\alpha_1}$ and $\psi_{2n-2} = 2\sigma_{-\alpha_1}$. The Chern classes c_1 and c_2 are also related to Schubert classes:

$$\begin{aligned} c_1 &= [\{V_2 \in \mathbb{G}_Q(2, 2n+1) \mid \dim(W_2^\perp \cap V_2) \geq 1\}] \\ c_2 &= [\{V_2 \in \mathbb{G}_Q(2, 2n+1) \mid V_2 \subset W_1^\perp\}] \end{aligned}$$

or with the notation of Section 2: $c_1 = \sigma_{\Theta - \alpha_2}$ and $c_2 = \sigma_{\Theta - \alpha_2 - \alpha_1}$.

To compute quantum cohomology, we need to deform the preceding relations using the quantum parameter q . Remark that the degree of q is $2n - 2$ therefore only the last two equations in the presentation (4) can be deformed. We thus need to compute the quantum terms in the equations (5). Now using Example 4.4, we can easily compute the q terms in the first equation of (5). We get

$$\psi_{2n-2} + c_1 * \psi_{2n-3} + c_2 * \psi_{2n-4} = 2q. \quad (6)$$

To deform the last equation, we only need to use the affine symmetries (see Subsection 2.5). Indeed $\frac{1}{2}\psi_{2n-2}$ corresponds to the opposite of the cominuscul simple root. We get

$$c_2 * \psi_{2n-2} = 2qc_2. \quad (7)$$

To study the semi-simplicity of the quantum cohomology ring, we only need to prove that when we set $q = 1$, the scheme $\text{Spec}(QH^*(\mathbb{G}_Q(2, 2n+1), \mathbb{Q})_{q=1})$ is reduced. In turn, since $QH^*(\mathbb{G}_Q(2, 2n+1), \mathbb{Q})$ is the ring of S_2 -invariants of the quotient $\mathbb{Q}[x_1, x_2]/(\psi_{2n-2} + c_1 * \psi_{2n-3} + c_2 * \psi_{2n-4} - 2q, c_2 * \psi_{2n-2} - 2qc_2)$, it is enough to check that this quotient is reduced.

For this we solve equations (6) and (7). We get

$$-\sum_{k=0}^{n-1} x_1^{2k} x_2^{2n-2-2k} = 2q \quad \text{and} \quad x_1^2 x_2^2 \sum_{k=0}^{n-2} x_1^{2k} x_2^{2n-4-2k} = 2qx_1 x_2.$$

To solve these equations for $q = 1$, we first remove the solutions (x_1, x_2) with $x_i = 0$ and x_{3-i} an $2(n-1)$ -th root of -2 . There are $4(n-1)$ distinct such solutions. Then we assume $x_1 x_2 \neq 0$ and write $x_2 = \lambda x_1$. We get

$$-x_1^{2(n-1)}(1 + \lambda^2 + \dots + \lambda^{2(n-1)}) = 2 \quad \text{and} \quad x_1^{2(n-1)}\lambda(1 + \lambda^2 + \dots + \lambda^{2(n-2)}) = 2.$$

Equating these relations and dividing by $x_1^{2(n-1)}$ we get:

$$1 + \lambda + \dots + \lambda^{2n-2} = 0$$

therefore λ is a $(2n-1)$ -th root of unity different from 1. Remark that therefore $\lambda \neq -1$ and we have

$$1 + \lambda^2 + \dots + \lambda^{2(n-1)} = \frac{\lambda^{2n} - 1}{\lambda^2 - 1} = \frac{\lambda - 1}{\lambda^2 - 1} = \frac{1}{\lambda + 1}.$$

There are $2n-2$ different solutions for λ and then x_1 is a $(2n-2)$ -th root of $-2(\lambda+1)$ and there are $2n-2$ solutions for x_1 . We described $4(n-1) + (2n-2)(2n-2) = 2n(2n-2)$ solutions therefore the above affine scheme is reduced. $\square$

Proposition 6.5 *The localised quantum cohomology algebra $QH^*(F_4/P_1, \mathbb{Q})_{loc}$ is semi-simple. It therefore admits an algebra involution sending q to its inverse.*

Proof. Recall that we have a presentation of the quantum cohomology ring given by

$$QH^*(F_4/P_1, \mathbb{C}) = \mathbb{C}[h, s, q]/(-h^8 + 12s^2 + 16q, 3h^{12} - 18h^8s + 24h^4s^2 + 8s^3)$$

where h is of degree 1 and s of degree 4. Recall also that $c_1 = 8$. We solve these equations in h and s when setting $q = 1$. First remark that there is no solution on the closed subset $s = 0$ neither on the closed subset $h = 0$. Eliminating h^8 in the second equation using the first one we get

$$3h^4(5s^2 + 4) = 4s(13s^2 + 18).$$

Taking the square of this equation and eliminating h^8 again we get

$$s^6 - 108s^4 - 576s^2 - 576 = 0. \tag{8}$$

For $P(S) = S^3 - 108S^2 - 576S - 576$ we have $P(-2) = 136$ and $P(0) = -576$, thus (8) is a degree 3 equation in s^2 whose solutions are real and distinct. The result follows. $\square$

Proposition 6.6 *The localised quantum cohomology algebra $QH^*(G_2/P_2, \mathbb{Q})_{loc}$ is semi-simple. It therefore admits an algebra involution sending q to its inverse.*

Proof. Recall that we have a presentation of the quantum cohomology ring given by

$$QH^*(G_2/P_2, \mathbb{C}) = \mathbb{C}[h, q]/(h^6 - 18h^3q - 27q^2)$$

where h is of degree 1. Recall also that $c_1 = 3$. We solve this equation in h when setting $q = 1$. This is a degree 2 equation in h^3 whose solutions are real non vanishing and distinct. $\square$

6.3 Some properties of the involution for adjoint non coadjoint varieties

Recall that in the minuscule and cominuscule cases, it was proved in [ChMaPe07] and [ChMaPe09a] that the involution ι given by complex conjugation has the nice property of sending a Schubert class to a multiple of such a class. In this section we shall see that contrary to these cases this property is not satisfied.

We shall first prove a nice property of the involution ι , namely it sends a class to a multiple of itself and the quantum parameter q for classes of degree c_1 .

Proposition 6.7 *For X an adjoint non coadjoint rational homogeneous space, let σ be a Schubert class of degree dc_1 for some integer d , then we have*

$$\iota(\sigma) = \frac{\sigma}{q^{2d}}$$

Proof. To prove this result, we may restrict ourselves to the specialisation of the quantum cohomology obtained by setting $q = 1$. The classes of this specialisation can be seen as functions with values in $\mathbb{C}$ on the variety $\text{Spec}(QH^*(X, \mathbb{C})_{q=1})$. The involution ι is given by complex conjugation on these functions. It is thus enough to prove that the values of these functions for classes of degree multiple by c_1 are real.

For this we prove that σ annihilates a polynomial equation whose solutions are real. This polynomial equation will be the minimal polynomial of the endomorphism given by the multiplication by σ . Let us first remark that it is enough to compute the minimal polynomial of the multiplication by σ on the subspace of degree c_1 classes. Indeed, assume that $P(\sigma)$ vanishes on that subspace, then $P(\sigma)q = 0$ and therefore because we inverted q we have $P(\sigma) = 0$.

We therefore restrict ourselves on the subspace H of degree c_1 classes and prove that the endomorphism given by multiplication by σ is adjoint for a non degenerate real quadratic form. This will prove the result because the eigenvalues of such endomorphisms are real.

Let us define a pairing on H by

$$(a, b) = \text{coefficient of } q^2 \text{ in the product } a * b.$$

This pairing is clearly symmetric and we shall prove that the quantum Schubert classes of degree c_1 form an orthogonal basis for this pairing. Indeed, denote by pt the class of a point in X , this orthogonality result comes from a formula for the multiplication by the class pt . This formula is a consequence of a more general formula for the multiplication by the Schubert class corresponding to the reflection with respect to the highest root (see [LaSh09, Proposition 11.2]) which is pt in our situation:

Fact 6.8 *We have, for α a long simple root and ϖ_α the associated fundamental weight, the formula:*

$$\text{pt} * \sigma_{-\alpha} = \langle \varpi_\alpha, \Theta^\vee \rangle q^2 \sigma_\alpha$$

In particular the quantum Schubert classes in H form an orthogonal basis and the form is positive definite. But now we may compute

$$(\sigma * a, b) = \text{coefficient of } q^2 \text{ in the product } \sigma * a * b = (a, \sigma * b)$$

and σ is adjoint from the form $(\ , \)$. This concludes the proof. $\square$

However, we will now prove that the involution ι is not as simple as in the minuscule or cominuscule cases. In particular, we prove the following

Theorem 6 *For X an adjoint non coadjoint rational homogeneous space, there exists a Schubert class σ such that $\iota(\sigma)$ is not a multiple of a Schubert class.*

Proof. Remark that to prove this result, we may set $q = 1$ so that the involution is the identity on degree c_1 classes. The proof will be a case by case analysis even if the idea of the proof is always the same: take σ a class of degree a divisor of c_1 , say $d \deg(\sigma) = c_1$. Then we have $\deg(\sigma^d) = c_1$ therefore $\iota(\sigma)^d = \iota(\sigma^d) = \sigma^d$ by Proposition 6.7. We are then reduced to prove that σ is the only Schubert class τ such that $\deg(\tau) \equiv -\deg(\sigma) \pmod{c_1}$ and τ^d is a scalar multiple of σ^d . When $\deg(\sigma) \equiv -\deg(\sigma) \pmod{c_1}$ then we will also need to prove that $\iota(\sigma)$ is not a scalar multiple of σ .

Let us start with $X = G_2/P_2$. Recall that in this situation $c_1 = 3$. Take $\sigma = h$. We have by Chevalley formula the equality $\sigma^3 = 6\sigma_3 + 3q$ where σ_3 is the only Schubert class of degree 3. If the class $\iota(\sigma)$ is a multiple of a Schubert class, then it has to be a multiple of σ_2 the unique degree 2 Schubert class or a multiple of pt the class of a point. But $3\sigma_2 = h^2$ therefore we may compute σ_2^3 using the equation of the presentation of the cohomology ring 5.1 we get $3\sigma_2^3 = 2h^3q + 3q^2$ and by Chevalley formula $\sigma_2^3 = 4\sigma_3q + 3q^2$ which is not a scalar multiple of $\iota(\sigma^3) = 6\sigma_3 + 3q$. We can also compute $\text{pt}^3 = 2q^2\ell * \text{pt}$ where ℓ is the class of a line in X (see Corollary 3.23). But by Theorem 3 we have $\text{pt} * \ell = \lambda q^2\sigma_3$ because the other term would be q^3 (by [LaSh09, Proposition 11.2] or several applications of the Chevalley formula we even have $\text{pt} * \ell = q^2\sigma_3$) therefore $\text{pt}^3 = 2\lambda q^4\sigma_3$ which is not a scalar multiple of $\iota(\sigma^3) = 6\sigma_3 + 3q$.

We now deal with $X = F_4/P_1$. Recall that in that situation $c_1 = 8$. Take σ to be the Schubert class associated to the element $s_4s_3s_2s_1$ in the Weyl group. We have $\sigma^2 = \sigma_{-\alpha_1} + \sigma_{-\alpha_2}$ where α_1 and α_2 are simple root. This formula can be proved using the presentation of Proposition 5.2, we proved it using our program [Ch09]. If the class $\iota(\sigma)$ is a multiple of a Schubert class, then it has to be a multiple of σ , of τ the other degree 4 Schubert class or of σ_{12} the unique degree 12 Schubert class. Using our program, we have $\tau^2 = 6\sigma_{-\alpha_1} + 8\sigma_{-\alpha_2}$ and $\sigma_{12}^2 = 3q^2\sigma_{-\alpha_1} + 5q^2\sigma_{-\alpha_2} + q^3$. Therefore none of them is a multiple of $\iota(\sigma)^3 = \sigma_{-\alpha_1} + \sigma_{-\alpha_2}$. To conclude, we need to prove that $\iota(\sigma)$ is not equal to a multiple of σ . For this we consider $\iota(\sigma)$ as a function on the scheme $\text{Spec}(QH^*(X, \mathbb{C})_{q=1})$. In the basis $(q, \sigma_{-\alpha_1}, \sigma_{-\alpha_2})$ of H the matrix of the multiplication by $\iota(\sigma)^2 = \sigma^2$ is given by (once more we use our program):

$$\begin{pmatrix} 0 & 2 & 3 \\ 1 & 5 & 9 \\ 1 & 6 & 10 \end{pmatrix}$$

whose minimal polynomial is $P(x) = x^3 + 15x^2 + 9x + 1$. It has two real positive roots and one real negative root, and from $P(\sigma^2) = 0$ we deduce that σ has four real values and two complex conjugate values. Its complex conjugate is therefore not a multiple of itself.

We finish with $X = \mathbb{G}_Q(2, 2n+1)$. Recall that in that situation $c_1 = 2n-2$. We first prove that there is a Schubert class σ such that its square σ^2 is a multiple of $\sigma_{-\alpha_{n-1}}$. For this we need to discuss on the parity of n . We take for σ the Schubert class associated to the Schubert variety (here we fix a complete isotropic flag $(W_i)_{i \in [1, n]}$):

$$\begin{cases} \{V_2 \in \mathbb{G}_Q(2, 2n+1) / V_2 \subset W_p^\perp\} & \text{if } n-1 = 2p \text{ is even,} \\ \{V_2 \in \mathbb{G}_Q(2, 2n+1) / V_2 \subset W_{p+2}\} & \text{if } n-1 = 2p+1 \text{ is odd.} \end{cases}$$

Remark that the degree of σ is given by

$$\deg(\sigma) = \begin{cases} n-1 & \text{if } n-1 \text{ is even,} \\ 3(n-1) & \text{if } n-1 \text{ is odd.} \end{cases}$$

Fact 6.9 *The square σ^2 is a scalar (and q) multiple of $\sigma_{\alpha_{n-1}}$.*

Proof. Let us deal with the case $n - 1$ even first. By Example 4.4, there is no q term in σ^2 . This is of degree $2(n - 1)$ and we need to compute the classical products $\sigma \cup \sigma \cup \sigma_{\alpha_k}$ for α_k a simple root and $k \in [1, n - 1]$. The Schubert variety associated to the class σ_{α_k} is

$$\{V_2 \in \mathbb{G}_Q(2, 2n + 1) / V_2 \subset W_{k-1}^\perp \text{ and } \dim(V_2 \cap W_{k+1}) \geq 1\}$$

where $(W_i)_{i \in [1, n]}$ is a complete isotropic flag. We therefore want to count the number of elements $V_2 \in \mathbb{G}_Q(2, 2n + 1)$ such that

$$V_2 \subset W_p^\perp, V_2 \subset T_p^\perp, V_2 \subset U_{k-1}^\perp \text{ and } \dim(V_2 \cap U_{k+1}) \geq 1$$

where $(U_i)_{i \in [1, n]}$, $(T_i)_{i \in [1, n]}$ and $(W_i)_{i \in [1, n]}$ are three generic complete isotropic flags. We see that these conditions imply that $\dim(U_{k+1} \cap W_p^\perp \cap T_p^\perp) \geq 1$ and because these flags are in general position that $k + 1 - 2p \geq 1$ therefore $k \geq n - 1$. In particular the only Schubert class appearing in σ^2 is $\sigma_{-\alpha_{n-1}}$ and the result follows. We even know (from [FuWo04]) that this product is not zero and easy manipulations with isotropic subspaces show that $\sigma^2 = 2\sigma_{-\alpha_{n-1}}$ (we will not need this fact here).

Now assume $n - 1$ to be odd. By Corollary 3, there is no q^3 term in σ^2 , and by Example 4.4, there is no q term, thus there are only terms of the form $q^2\sigma_{-\alpha_k}$ for α_k a simple root and $k \in [1, n - 1]$. We therefore need to compute $I_2(\sigma, \sigma, \sigma_{\alpha_k})$. But if $f : \mathbb{P}^1 \rightarrow \mathbb{G}_Q(2, 2n + 1)$ is a degree 2 morphism then all the subspaces in the image are contained in a 4-dimensional, non isotropic in general, subspace V_4 . We shall use this subspace to get some information. Indeed, if f meets three general Schubert varieties representing the Schubert classes σ , σ and σ_{α_k} then the subspace V_4 satisfies:

$$\begin{aligned} \dim(V_4 \cap W_{p+2}) &\geq 2, \\ \dim(V_4 \cap T_{p+2}) &\geq 2, \\ V_4 \cap U_{k-1}^\perp &\text{ contains a 2-dimensional isotropic subspace and} \\ \dim(V_4 \cap U_{k+1}) &\geq 1, \end{aligned}$$

where $(U_i)_{i \in [1, n]}$, $(T_i)_{i \in [1, n]}$ and $(W_i)_{i \in [1, n]}$ are three generic complete isotropic flags. We see that these conditions imply that $\dim(U_{k+1} \cap (W_{p+2} + T_{p+2})) \geq 1$ and because these flags are in general position that $k + 1 + p + 2 + p + 2 - (2n + 1) \geq 1$ therefore $k \geq n - 1$. We conclude as before. Again one can prove the equality $\sigma^2 = 2q^2\sigma_{-\alpha_{n-1}}$. $\square$

We now prove that there is no Schubert class τ different from σ with $\deg(\tau) \equiv -\deg(\sigma) \equiv n - 1 \pmod{c_1}$ and τ^2 equal to a scalar multiple of $\sigma_{-\alpha_{n-1}}$. For this we describe the Schubert varieties associated to such classes τ of degree equal to $n - 1$ modulo c_1 :

$$\begin{aligned} \{V_2 \in \mathbb{G}_Q(2, 2n + 1) / V_2 \subset W_a^\perp \text{ and } \dim(V_2 \cap W_{a+b+1}) \geq 1\} &\text{ with } a \in [0, \frac{n-1}{2}[\text{ and } 2a + b = n - 1 \\ \{V_2 \in \mathbb{G}_Q(2, 2n + 1) / V_2 \subset W_{a+b+2} \text{ and } \dim(V_2 \cap W_{a+1}) \geq 1\} &\text{ with } a \in [0, \frac{n-2}{2}[\text{ and } 2a + b = n - 2 \end{aligned}$$

where $(W_i)_{i \in [1, n]}$ is a complete isotropic flag. We want to compute the coefficient of $\sigma_{-\alpha_{n-1}}$ (resp. $q^2\sigma_{-\alpha_{n-1}}$) is the square τ^2 in the first (resp. second) example.

Fact 6.10 *The only Schubert class τ of degree $n - 1$ or $3(n - 1)$ such that τ^2 is a multiple of σ^2 by a scalar product of a power of q are the classes $\pm\sigma$.*

Proof. For τ a degree $n - 1$ class, we proceed as follows: first we prove that with notation as in Proposition 6.4, we have $\sigma_2^{n-1} = \sigma^2$ (recall that σ_2 is the degree 2 cohomology class of the Schubert variety $\{V_2 \in \mathbb{G}_Q(2, 2n + 1) / V_2 \subset W_1^\perp\}$ with $(W_i)_{i \in [1, n]}$ is a complete isotropic flag). Indeed, we can easily compute using jeu de taquin (cf. [ChPe09a]) that $\sigma_2^k = \sigma_{2k}$ where σ_{2k} is the cohomology class of the Schubert variety $\{V_2 \in \mathbb{G}_Q(2, 2n + 1) / V_2 \subset W_k^\perp\}$ with $(W_i)_{i \in [1, n]}$ is a complete isotropic flag

and $k \leq n - 2$. Then we easily see (use for example the heaps like in [ChPe09a]) that the only degree $2n - 2$ class containing σ_{n-1} is $\frac{1}{2}\sigma^2$ and from Example 4.4 there is no q term in the product $\sigma_2 * \sigma_{n-2}$. Therefore σ_2^{n-1} is a scalar multiple of $\frac{1}{2}\sigma^2$. From this one easily get $\sigma_2^{n-1} = \sigma^2$ (we will only need that it is a scalar multiple of σ^2).

Recall from our presentation of the quantum cohomology in Proposition 6.4 that any degree $n - 1$ class τ can be expressed as a symmetric degree $n - 1$ polynomial in the variables x_1 and x_2 . Moreover, the first equation being in degree $2n - 2$ and of the form $q = P(x_1, x_2)$ with P a degree $2n - 2$ polynomial, we have that, in the subring generated by x_1 and x_2 in $QH^*(\mathbb{G}_Q(2, 2n + 1), \mathbb{C})$, the multiplication is injective from degree $n - 1$ classes to degree $2n - 2$ classes. In particular, the only classes τ such that $\tau^2 = \sigma^2 = \sigma_2^{n-1}$ is $\pm\sigma_2^p$ with $2p = n - 1$ for $n - 1$ even and no class for $n - 1$ odd. Now $\sigma_2^p = \sigma_{2p}$ as above and is the class σ .

For the second case, we were not able to do the same trick and we need to do geometry on subspaces as in Fact 6.9. We need to compute the square of the class τ of a Schubert variety $\{V_2 \in \mathbb{G}_Q(2, 2n + 1) / V_2 \subset W_{a+b+2} \text{ and } \dim(V_2 \cap W_{a+1}) \geq 1\}$ with $a \in [0, \frac{n-2}{2}[$ and $2a + b = n - 2$ and where $(W_i)_{i \in [1, n]}$ is a complete isotropic flag. To prove that its square is not a scalar multiple of $q^2\sigma^2$, we compute its coefficient on the class $\sigma_{-\alpha_{n-b-1}}$. We therefore have to compute the Gromov-Witten invariant $I_2(\tau, \tau, \sigma_{\alpha_{n-b-1}})$. For this we want to count the number of 4-dimensional (non isotropic) subspaces V_4 such that

$$\begin{aligned} \dim(V_4 \cap W_{a+1}) &\geq 1, & \dim(V_4 \cap W_{a+b+2}) &\geq 2, \\ \dim(V_4 \cap T_{a+1}) &\geq 1, & \dim(V_4 \cap T_{a+b+2}) &\geq 2, \\ \dim(V_4 \cap U_{n-b}) &\geq 1, & V_4 \cap U_{n-b-2}^\perp &\text{ contains a 2-dimensional isotropic subspace,} \end{aligned}$$

where $(U_i)_{i \in [1, n]}$, $(T_i)_{i \in [1, n]}$ and $(W_i)_{i \in [1, n]}$ are three generic complete isotropic flags.

Suppose we are given such V_4 , and let us analyse some properties of this linear subspace. Since $(U_i)_{i \in [1, n]}$, $(T_i)_{i \in [1, n]}$ and $(W_i)_{i \in [1, n]}$ are generic, the above inequalities are equalities. We denote P the intersection point of $\mathbb{P}V_4$ and $\mathbb{P}U_{n-b}$. We denote $L_1 = V_4 \cap W_{a+b+2}$ and $L_2 = V_4 \cap T_{a+b+2}$. In the plane $\mathbb{P}(V_4 \cap P^\perp)$ there are two isotropic lines through P , the line $\mathbb{P}L$ not meeting $\mathbb{P}L_1$ and $\mathbb{P}L_2$ and the line $\mathbb{P}L_3$. In the plane $\mathbb{P}\pi_P(V_4)$ we denote $p = \mathbb{P}\pi_P(L)$. In $\mathbb{P}\pi_L(L_4)$ we denote Λ_3 the projection of $\mathbb{P}L_3$. Note that $\pi_L(V_4) = \pi_L(V_4 \cap W_{a+1}) \oplus \pi_L(V_4 \cap T_{a+1})$ and therefore $\Lambda_3 \in \mathbb{P}(\pi_L(W_{a+1}) \oplus \pi_L(T_{a+1}))$. Finally we denote π_3 the projection with center generated by L and L_3 and we observe that $\Lambda := \mathbb{P}\pi_3(V_4) \in \pi_3(W_{a+1}) \cap \pi_3(T_{a+1})$.

We now explain how to find exactly two subspaces V_4 satisfying the above conditions given the isotropic flags.

First of all we see that $U_{n-b} \cap (W_{a+b+2} + T_{a+b+2})$ has dimension $n - b + a + b + 2 + a + b + 2 - (2n + 1) = 1$. Thus the point P is determined by $P = \mathbb{P}(U_{n-b} \cap (W_{a+b+2} + T_{a+b+2}))$. We then have $\dim \pi_P(W_{a+b+2}) = \dim \pi_P(T_{a+b+2}) = a + b + 2$ whereas $\dim \pi_P(W_{a+b+2} + T_{a+b+2}) = 2(a + b + 2) - 1$, thus the spaces $\mathbb{P}\pi_P(W_{a+b+2})$ and $\mathbb{P}\pi_P(T_{a+b+2})$ intersect in one point. Note moreover that $\pi_P(W_{a+b+2}) \supset \pi_P(L_1) \supset \pi_P(L)$, and similarly for $\pi_P(T_{a+b+2})$, and thus p is determined by $p = \mathbb{P}(\pi_P(W_{a+b+2}) \cap \pi_P(T_{a+b+2}))$. This defines uniquely the line $\mathbb{P}L$ in $\mathbb{P}\mathbb{C}^{2n+1}$.

Now we observe that $\mathbb{P}(U_{n-b-2}^\perp \cap P^\perp)$ meets $\mathbb{P}L$ at P (except for $(a, b) = (0, n - 2)$), so that $U_{n+b+1} := \pi_L(U_{n-b-2}^\perp \cap P^\perp)$ is a quadratic space of dimension $n + b + 1$ (for $(a, b) = (0, n - 2)$ the space $\pi_L(U_{n-b-2}^\perp \cap P^\perp)$ is of dimension $2n - 2$). The intersection $U_{n+b+1} \cap (\pi_L(W_{a+1}) + \pi_L(T_{a+1}))$ has dimension 2 and is non-degenerate for this quadratic form (for $(a, b) = (0, n - 2)$, the space $\pi_L(W_{a+1}) + \pi_L(T_{a+1})$ is already of dimension 2 with a non degenerate quadratic form). Since we have seen that $\Lambda_3 \in \mathbb{P}(\pi_L(W_{a+1}) \oplus \pi_L(T_{a+1}))$, it follows that we have exactly two choices for Λ_3 : the two isotropic points of $\mathbb{P}(U_{n+b+1} \cap (\pi_L(W_{a+1}) + \pi_L(T_{a+1})))$. Let us choose one isotropic point Λ_3 . This defines the projection π_3 . Observe that $\pi_3(W_{a+1})$ and $\pi_3(T_{a+1})$ meet in dimension 1 in

$\pi_3(W_{a+1} \oplus T_{a+1})$ of dimension $2a+1$, and therefore Λ is defined by $\Lambda = \mathbb{P}(\pi_3(W_{a+1}) \cap \pi_3(T_{a+1}))$. This defines V_4 as the inverse image of Λ by π_3 . $\square$

To conclude the proof we need to prove the following fact:

Fact 6.11 *We have $\iota(\sigma) \neq \pm\sigma$.*

Proof. For this we first specialise to $q = 1$ and we will prove that $\sigma^2 = 2\sigma_{-\alpha_{n-1}}$ takes positive and negative real values. This will imply that for its square root the complex conjugation (i.e. the involution ι) is never multiplication by ± 1 .

Consider the multiplication by $\sigma_{-\alpha_{n-1}}$ as an endomorphism u of the space H (the vector space generated by the degree $2n-2$ classes) and adjoint for the scalar product $(\ , \)$ defined in the proof of Proposition 6.7. First of all because the quantum multiplication of Schubert classes has non negative integer coefficients, we see that $\text{tr}(u) > 0$ and therefore u has positive eigenvalues. If all eigenvalues of u were non negative, then we would have $(u(x), x) \geq 0$ for all $x \in H$ with equality if and only if $u(x) = 0$. But take $x = 1 + \sigma_{-\alpha_1}$ (recall that multiplication by $\sigma_{-\alpha_1}$ induces the affine symmetry described in Subsection 2.5). We have $u(x) = 2\sigma_{-\alpha_{n-1}}$ which is non trivial but $(u(x), x) = 0$. Therefore U has negative eigenvalues and the result follows. $\square$

This concludes the proof: $\iota(\sigma)$ is never a Schubert class. $\square$

Remark 6.12 Let us compute explicitly in an example the element $\iota(\sigma)$ with σ as in the previous proposition. Assume that $X = \mathbb{G}_Q(2, 7)$. We have $c_1 = 4$. In that case, we have $\sigma = \sigma_2$ with the notation of Proposition 6.4. Recall also that there is a unique other Schubert class of degree 2 denoted τ_2 . We have $\sigma^2 = 2\sigma_{-\alpha_{ad}}$ and easy computations give the following equation:

$$s^2(\sigma^4 - 4\sigma^2 - 16) = 0.$$

Therefore the values of σ^2 are 0, $2(1 + \sqrt{5})$ and $2(1 - \sqrt{5})$. In particular we see that the polynomial

$$Q(x) = \frac{x(x^2 - 2)}{2\sqrt{5}}$$

maps the square roots of the values of σ^2 to their complex conjugate therefore

$$\iota(\sigma) = \frac{s^3 - 2sq}{2q^2\sqrt{5}} = \frac{1}{q^2\sqrt{5}}(2\ell + 2\tau_2 - \sigma_2)$$

where ℓ is the class of a line.

7 The incidence variety

In this section we deal with the adjoint variety X of type A_n , namely the incidence variety between points and hyperplanes in $\mathbb{P}^n$. For many aspects the results proved in the previous sections are still valid. However, as the Picard group of X is $\mathbb{Z}^2$, some of the results change. We will explain in this section which of the results proved remain valid and which changes are required for the others.

Fact 7.1 *(i) The parametrisation of Schubert classes by roots in Section 2 remains valid. In particular Proposition 2.8 is valid.*

(ii) As the Picard group is $\mathbb{Z}^2$, there are two classes of degree 1 that we shall denote by h_1 and h_2 and there are two quantum parameters q_1 and q_2 . Therefore, the parametrisation of quantum monomials by roots of the affine root system fails.

- (iii) Chevalley and Quantum Chevalley formulas also fail. However, setting $h = h_1 + h_2$ and $q = q_1 + q_2$, then the formula in Theorem 2 is true.
- (iv) The formula for affine symmetries fails by lack of a description of the quantum monomials.
- (v) The description of ϖ -minuscule elements is valid as well as the fact that $d_{\max} = 2$.

It is easy to derive an explicit formula for the quantum Chevalley formula. This formula completely determines the quantum cohomology ring as the degree 1 classes are generators. We get the following:

Proposition 7.2 *The quantum cohomology ring $QH^*(X, \mathbb{Q})$ is generated by h_1 , h_2 and the quantum parameters q_1 and q_2 . It is the quotient of $\mathbb{Q}[h_1, h_2, q_1, q_2]$ by the relations*

$$\sum_{k=0}^n h_1^k (-h_2)^{n-k} = q_1 + (-1)^n q_2 \text{ and } h_1^{n+1} = q_1 (h_1 + h_2).$$

- (iii) *The quantum cohomology ring is semi-simple for quantum parameters q_1 and q_2 satisfying*

$$q_1 \neq 0, \quad q_2 \neq 0 \text{ and } q_1 + (-1)^n q_2 \neq 0.$$

Proof. The first two formulas are direct interpretations of the quantum Chevalley formula given in [FuWo04].

Let us prove the semi-simplicity. By multiplication of the quantum parameters q_1 and q_2 by the same scalar, we may assume $q_1 + (-1)^n q_2 = 1$. Let us first remark that because $q_1 \neq 0$, if $h_1 = 0$ then from the second equation we have $h_2 = 0$. This is not compatible with the first equation. We may thus assume $h_1 \neq 0$ and set $h_2 = \lambda h_1$. We obtain

$$h_1^n \sum_{k=0}^n (-\lambda)^k = 1 \text{ and } h_1^n = q_1 (1 + \lambda).$$

Remark that because $h_1 \neq 0$ we have $1 + \lambda \neq 0$ therefore, multiplying the first equation by $q_1 (1 + \lambda)$ and equating, we get $h_1^n q_1 (1 + (-1)^n \lambda^{n+1}) = h_1^n$ and the equation

$$q_1 \lambda^{n+1} = (-1)^{n+1} (q_1 - 1).$$

Because $q_1 \neq 0$ (otherwise $h_1 = 0$) and $q_1 \neq 1$ (otherwise $q_2 = 0$) we get $n + 1$ distinct solutions for λ and $n(n + 1)$ solutions for (h_1, h_2) . $\square$

References

- [BeGeGe73] Bernšteĭn, I. N., Gel'fand, I. M., Gel'fand, S. I., *Schubert cells and the cohomology of a flag space*. Funkcional. Anal. i Priložen. **7** (1973), no. 1.
- [Bor53] Borel, A., *Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts*. Ann. of Math. (2) **57**, (1953).
- [Bou54] Bourbaki, N., *Groupes et algèbres de Lie*. Hermann 1954.
- [BrKu05] Brion, M., Kumar, S., *Frobenius splitting methods in geometry and representation theory*. Progress in Mathematics, 231. Birkhäuser Boston, Inc., Boston, MA, 2005.
- [BuKrTa03] Buch, A., Kresch, A., Tamvakis, H., *Gromov-Witten invariants on Grassmannians*. J. Amer. Math. Soc. **16** (2003), no. 4.

- [BuKrTa08] Buch, A., Kresch, A., Tamvakis, H., *Quantum Pieri rules for isotropic Grassmannians*. Preprint arXiv:0809.4966 (2008).
- [ChMaPe07] Chaput, P.-E., Manivel, L., Perrin, N., *Quantum cohomology of minuscule homogeneous spaces II : Hidden symmetries*. Int. Math. Res. Not. IMRN (2007), no. 22.
- [ChMaPe08] Chaput, P.-E., Manivel, L., Perrin, N., *Quantum cohomology of minuscule homogeneous spaces*. Transform. Groups **13** (2008), no. 1, 47–89.
- [ChMaPe09a] Chaput, P.-E., Manivel, L., Perrin, N., *Quantum cohomology of minuscule homogeneous spaces III : semi-simplicity and consequences*. To appear in Canadian J. of Math. (2009).
- [ChMaPe09b] Chaput, P.-E., Manivel, L., Perrin, N., *Affine symmetries of the equivariant quantum cohomology ring of rational homogeneous spaces*. To appear in Math. Res. Let. (2009).
- [ChPe09a] Chaput, P.-E., Perrin, N., *Towards a Littlewood-Richardson rule for Kac-Moody homogeneous spaces*. Preprint arXiv:0902.0152 (2009).
- [ChPe09b] Chaput, P.-E., Perrin, N., *Computations of quantum Giambelli formulas in some exceptional homogeneous spaces*. Preprint on the arXiv.
- [Ch09] Chaput, P.-E. *Java software for Schubert calculus*, available at www.math.sciences.univ-nantes.fr/~chaput/quiver-demo.html.
- [FuWo04] Fulton, W., Woodward, C., *On the quantum product of Schubert classes*. J. Algebraic Geom. **13** (2004), no. 4, 641–661.
- [Jac68] Jacobson, N., *Structure and representations of Jordan algebras*. American Mathematical Society Colloquium Publications, Vol. XXXIX American Mathematical Society, Providence, R.I. 1968.
- [Kac90] Kac, V. G., *Infinite-dimensional Lie algebras*. Third edition. Cambridge University Press, Cambridge, 1990.
- [KrTa03] Kresch, A., Tamvakis, H., *Quantum cohomology of the Lagrangian Grassmannian*. J. Algebraic Geom. **12** (2003), no. 4.
- [KrTa04] Kresch, A., Tamvakis, H., *Quantum cohomology of orthogonal Grassmannians*. Compos. Math. **140** (2004), no. 2.
- [Ful97] Fulton, W., *Young tableaux. With applications to representation theory and geometry*. London Mathematical Society Student Texts, 35. Cambridge University Press, Cambridge, 1997.
- [LaSh09] Lam, T., Shimizono, M., *Quantum cohomology of G/P and homology of affine Grassmannian*. To appear in Acta. Math. (2009).
- [LaMuSe79] Lakshmibai, V., Musili, C., Seshadri, C. S., *Geometry of G/P . III. Standard monomial theory for a quasi-minuscule P* . Proc. Indian Acad. Sci. Sect. A Math. Sci. **88** (1979), no. 3.
- [LaMa03] Landsberg, J. M., Manivel, L., *On the projective geometry of rational homogeneous varieties*. Comment. Math. Helv. **78** (2003), no. 1.

- [NiSe06] Nikolenko, S., Semenov N., *Chow ring structure made simple*. Preprint arXiv:math/0606335.
- [Pas08] Pasquier, B., *Fano horospherical varieties*. to appear in Bulletin de la SMF **136** (2008) no. 2.
- [Per02] Perrin, N., *Courbes rationnelles sur les variétés homogènes*. Ann. Inst. Fourier **52** (2002), no. 1.
- [Per07] Perrin, N., *Small resolutions of minuscule Schubert varieties*. Compos. Math. **143** (2007), no. 5.
- [Pro99a] Proctor, R. A., *Minuscule elements of Weyl groups, the numbers game, and d-complete posets*. J. Algebra **213** (1999), no. 1.
- [Ses78] Seshadri, C. S., *Geometry of G/P . I. Theory of standard monomials for minuscule representations*. C. P. Ramanujam—a tribute, Tata Inst. Fund. Res. Studies in Math., 8, Springer, Berlin-New York, 1978.
- [SiTi97] Siebert, B., Tian, G., *On quantum cohomology rings of Fano manifolds and a formula of Vafa and Intriligator*. Asian J. Math. **1** (1997), no. 4.
- [ThYo08] Thomas, H., Yong, A., *A combinatorial rule for (co)minuscule Schubert calculus*. To appear in Advances in Math.
- [Tho98] Thomsen, J. F., *Irreducibility of $\overline{M}_{0,n}(G/P, \beta)$* . Internat. J. Math. **9** (1998), no. 3.
- [Zak93] Zak, F. L., *Tangents and secants of algebraic varieties*. Translations of Mathematical Monographs, 127. American Mathematical Society, Providence, RI, 1993.

Pierre-Emmanuel CHAPUT,

Laboratoire de Mathématiques Jean Leray, UMR 6629 du CNRS, UFR Sciences et Techniques, 2 rue de la Houssinière, BP 92208, 44322 Nantes cedex 03, France and

Hausdorff Center for Mathematics, Universität Bonn, Villa Maria, Endenicher Allee 62, 53115 Bonn, Germany

email: pierre-emmanuel.chaput@math.univ-nantes.fr.

Nicolas PERRIN,

Hausdorff Center for Mathematics, Universität Bonn, Villa Maria, Endenicher Allee 62, 53115 Bonn, Germany and

Institut de Mathématiques de Jussieu, Université Pierre et Marie Curie, Case 247, 4 place Jussieu, 75252 Paris Cedex 05, France.

email: nicolas.perrin@hcm.uni-bonn.de.